

2012

UF

IFAS

UNIVERSITY of FLORIDA

University of Florida's Pest Control Guide for Turfgrass Managers



2012 University of Florida's Pest Control Guide for Turfgrass Managers

Compiled and Edited by Dr. J. Bryan Unruh
Professor and Extension Turfgrass Specialist

This guide supplies information concerning pesticides that can be used for pests in turfgrasses. Because of the risks involved in their handling and use, the U. S. Environmental Protection Agency has classified some pesticide products for restricted use. Such products are identified by the words "RESTRICTED USE PESTICIDE" placed above the product's brand name on the label. Such products must be applied by certified and licensed pesticide applicators or someone working under their direct supervision. Pesticide products that do not bear the "Restricted Use" designation can be purchased and applied by anyone. However, Florida law requires anyone who applies any pesticide to lawns associated with structures (residences, commercial buildings, etc.) for monetary compensation to be licensed or supervised by someone with a license. Persons who apply pesticides to golf courses, parks, cemeteries, and athletic fields must be licensed or supervised by someone with a license only if a restricted use pesticide is applied. Government employees who apply pesticides to the lawns around government buildings and private business property owners who apply pesticides to the lawns around the buildings on their business property must also be licensed. The Florida Department of Agriculture and Consumer Services licenses pesticide applicators. See Pesticide Licensing Category Information (page 7) for specific information on types of licenses required for application of pesticides to turf/lawns in Florida.

Use pesticides safely to protect against human injury and harm to the environment. Diagnose your pest problem; select the proper pesticide, if one is needed; follow the label directions; and obey all federal and state pesticide laws and regulations.

Use of brand names in this publication does not imply endorsement of the products or criticism of similar ones not mentioned, but are used herein for convenience only. Mention of a proprietary product does not constitute a guarantee or warranty of the product by the authors.

CONTENTS

<u>Title</u>	<u>Page</u>
<i>University of Florida Turfgrass Faculty</i>	2
<i>What's New in 2012</i>	3
<i>Turfgrass Publications</i>	6
<i>Pesticide Licensing Category Information</i>	7
<i>Insect Management</i>	11
<i>Disease Management</i>	18
<i>Nematode Management</i>	26
<i>Weed Management</i>	29
<i>Plant Growth Retardants Used in Turfgrass Management</i>	65
<i>Activated Charcoal for Pesticide Deactivation</i>	70
<i>Weed Control in Irrigation Water Supplies</i>	71
<i>Pesticide Calibration Formulas and Information.</i>	72
<i>Metric System Conversion Factors</i>	76
<i>Pesticide Application Record</i>	81
<i>Poison Control Center</i>	82

UNIVERSITY OF FLORIDA TURFGRASS FACULTY

Faculty Name	Phone Number	e-mail Address	Turf Specialty	Assignment Teaching/Research/Extension	% Turf
<i>Belle Glade</i>					
Dr. Ron Cherry	561/993-1529	pinesnpets@aol.com	Insects	0/100/0	40
<i>Ft. Lauderdale</i>					
Dr. John Cisar	954/577-6336	jlci@ufl.edu	Water, Turfgrass Management	0/60/40	100
Dr. Robin Giblin-Davis	954/577-6333	giblin@ufl.edu	Nematodes	0/100/0	50
<i>Gainesville</i>					
Dr. Eileen Buss	352/273-3976	eabuss@ufl.edu	Insects	5/25/70	75
Dr. Billy Crow	352/273-3941	wtr@ufl.edu	Nematodes	0/40/60	70
Dr. Jason Kruse	352/273-4569	jkk@ufl.edu	Turfgrass Management	70/30/0	100
Dr. Michael Dukes	352/392-1864 ext 205	MDDukes@ufl.edu	Ag. Engineering / Turf Irrigation	40/60/0	75
Dr. Howard Frank	352/273-3922	jhfrank@ufl.edu	Biological Control Mole Crickets	10/70/20	10
Dr. Philip Harmon	352/273-4622	pfharmon@ufl.edu	Diseases	0/40/60	33
Dr. George Hochmuth	352/392-1803 ext. 318	hoch@ufl.edu	Soil Fertility and Plant Nutrition	20/20/60	50
Dr. Kevin Kenworthy	352/392-6189	kenworth@ufl.edu	Plant Breeding	30/70/0	100
Dr. Jerry Sartain	352/392-7271	sartain@ufl.edu	Nutrition, Fertility	Emeritus	
Dr. Laurie Trenholm	352/273-4524	ltr@ufl.edu	Urban Landscape Turf Management	0/30/70	100
<i>Milton / Jay</i>					
Dr. Barry Brecke	850/995-3720 ext. 106	bjbe@ufl.edu	Weed Science	Emeritus	
Dr. Ramón León	850/995-3720 ext. 105	rglg@ufl.edu	Weed Science	0/60/40	50
Dr. Bryan Unruh	850/995-3720 ext. 108	jbu@ufl.edu	Culture, Turfgrass Management	70/0/30	100

What's New for 2012

Turfgrass Insect Control

1. **Acelepryn™** (DuPont) is now labeled for control of grubs, caterpillars, and billbugs. Chinch Bug suppression is also on the label.

Turfgrass Disease Control

1. Rapid Turfgrass Diagnostic Service now available. The service is offered through the Florida Extension Plant Disease Clinic in Gainesville. Details of the new service are available at <http://turf.ufl.edu>.

2. New Product Registrations:

Segway fungicide with the active ingredient cyazofamid (FMC) and Stellar fungicide with fluopicolide (Valent) have received registrations and include Pythium diseases on the labels. The products primarily have been evaluated on bentgrass, but warm-season turfgrass uses are not prohibited on the labels.

Disarm contains a QoI active fluoxastrobin (Arysta) and is available in several formulations, some premixed with other active ingredients.

Torque fungicide contains the newly registered DMI active ingredient tebuconazole (Cleary's). DMI actives have shown the potential to damage bermudagrass turf in Florida. Seashore paspalum and zoysiagrass are less likely to be damaged.

Turfgrass Nematode Control

1. **MustGro Invest:** Sometime in 2012 a new bionematicide made from formulated mustard material tradenamed MustGro Invest will be launched in Florida. This product is labeled for turfgrass and ornamental plant use.
2. UF has new thresholds for sting nematode on ultradwarf bermudagrass cultivars.

Turfgrass Weed Control

1. sulfentrazone + metsulfuron-methyl - **Blindside™** 66 WDG (FMC)

Rate: 6.5 to 10 oz. product/A

New postemergence herbicide for control of broadleaf weeds in bermudagrass, centipedegrass, St. Augustinegrass and zoysiagrass. Labeled for use on residential and commercial lawns, athletic fields, sod farms and golf course fairways and roughs.

2. thien carbazonone + iodosulfuron + dicamba - **Celsius™** 68 WDG (Bayer Environmental Science)

Rate: 2.5 to 4.9 oz. product/A

New postemergence herbicide for control of broadleaf weeds and some grasses in St. Augustinegrass, bermudagrass, centipedegrass, and zoysiagrass. This product is intended for use by commercial licensed applicators only. It is labeled for use on residential and commercial lawns, golf courses, sports fields, recreational areas, sod farms, roadsides and school grounds.

3. sulfentrazone + imazethapyr - **Dismiss® South** 4 SC (FMC)

Rate: 9.5 to 14.4 fl. oz./A

Postemergence control similar the that provided by Dismiss with added nutsedge control in

bahiagrass, bermudagrass, centipedegrass and zoysiagrass. It is labeled for use on residential and commercial lawns, athletic fields, sod farms, golf course fairways and roughs.

4. flazasulfuron - **Katana**[®] 25 WDG (PBI /Gordon)

Rate: 1.5 to 3 oz. product/A

New postemergence product for control of broadleaf weeds, sedges and some grass species and for removal of winter overseed cool-season grasses in bermudagrass, zoysiagrass and centipedegrass . Katana can be applied to turfgrass on golf course fairways and roughs, industrial parks, sod farms, cemeteries, professionally managed sports fields, and non-residential commercial lawns.

5. sulfentrazone + quinclorac - **Solitare**[®] 75 WDG (FMC)

Rate: 16 to 32 oz. product/A

This product provides postemergence control o grass, broadleaf weeds and sedges in bermudagrass, centipedegrass seashore paspalum and zoysiagrass. Solitare is labeled for use on residential and commercial lawns, athletic fields, sod farms and golf course fairways and roughs.

6. indaziflam - **Specticle** 20 WSP (Bayer Environmental Science)

Rate: 2.5 to 5.0 oz. product/A

Specticle provides preemergence control of annual grasses and broadleaf weeds in bermudagrass, zoysiagrass, centipedegrass, St. Augustinegrass, seashore paspalum and bahiagrass. It can be applied to turfgrass grown for residential and commercial lawns, golf course fairways, tees and roughs, sod, sports fields, parks and cemeteries.

7. carfentrazone + quinclorac - **SquareOne**[™] 70 WDG (FMC)

Rate: 12 to 18 oz. product/A

This product provides postemergence control of grass and broadleaf weeds in bermudagrass, centipedegrass, seashore paspalum and zoysiagrass. SquareOne is labeled for use on residential and commercial lawns, athletic fields, sod farms and golf course fairways and roughs.

8. dimethenamid - **Tower**[®] 6 EC (BASF)

Rate: 21 to 32 fl. oz./A

Tower provides preemergence control of annual grass and broadleaf weeds in bahiagrass, bermudagrass, centipedegrass, St. Augustinegrass, seashore paspalum and zoysiagrass. For use only on golf course fairways and roughs. Not for use on residential turf, commercial turf or sod.

9. **MSMA**

MSMA can no longer be **sold** in Florida for any turfgrass use

Under the original agreement between EPA and MSMA manufacturers MSMA use was to have ceased on December 31, 2010. However, the agreement was revised and MSMA can be used in Florida turfgrass until supply is depleted as long as the turfgrass use is on the label. Use must cease by December 31, 2013.

Plant Growth Regulators

1. Flurprimidol + Trinexapac-ethyl - **Legacy™** (SePRO)

Legacy Turf Growth Regulator (TGR) integrates patented synergistic turf growth regulator technology to reduce stem elongation and leaf blade length in perennial cool- and warm-season turfgrasses on golf courses resulting in a more compact and dense growth habit. Do not apply to bermudagrass putting greens or overseeded bermudagrass putting greens.

Turf Species: Tifway Bermudagrass, Seashore Paspalum
Rate: 10 to 20 fl oz/A

Turf Species: TifSport Bermudagrass, Zoysiagrass
Rate: 8 to 16 fl oz/A

Turf Species: Perennial Ryegrass overseeded bermudagrass
Rate: 15 to 30 fl oz/A

UNIVERSITY OF FLORIDA'S TURFGRASS PUBLICATION ORDER INFORMATION

Pests that Wreck Your Grass and Ruin Your Weekend! - SP 327

Did the bugs throw a party in your backyard and forget to invite you? Turn your fresh, green carpet of lawn into a patchy brown mat? Then it's time to learn about grass pests with Pests that wreck your grass and ruin your weekend! Use this colorful, informative booklet to get the lowdown on everything from armyworms to spittlebugs—before they get the better of you.

Designing, Construction, and Maintaining Bermudagrass Sports Fields - SP 361

This is the Second Edition of the definitive text on the science and practice of Bermudagrass sports fields: a must-have publication for those involved with designing, constructing and/or maintaining football or soccer fields, baseball and softball diamonds. Topics covered? Field drainage (both surface and subsurface); irrigation; turf establishment; "grow-in"; cultural practices; overseeding; pest control; preparations for special events; renovating damaged areas, etc. Spiral bound, 100 pages.

Florida Lawn Handbook - SP 45

Written in practical language by turfgrass experts, this new highly-anticipated edition is completely up-to-date, with the most current lawn management information. Color plates identify various grass types, weeds, diseases, and insects, including those that are good for your lawn! Chapters cover selection, adaptability, establishment, and maintenance for each type of lawn; soil analysis and fertilization; yearly calendars for lawn care and culture; mowing, watering, and calibrating sprinkler systems and fertilizer spreaders; overseeding for winter color; preparing a lawn for drought and low temperatures; weed and thatch control; safe pesticide application and use; the latest integrated pest management strategies; and complete, illustrated diagnostic information for weeds, diseases, insect problems, nematodes, and other pests. Whether you're an amateur or a pro, The Florida Lawn Handbook is an invaluable aid to growing a beautiful, healthy lawn year round.

Ornamental and Turf Pest Control (Ornamental and Turf; Lawn and Ornamental Exams) - SM 007

This is the exam preparation and general reference manual for commercial or public applicators seeking certification and licensure to apply pesticides for ornamental and turf pest control. The information includes weeds, insects, diseases, and nematodes affecting ornamental plants and turf and their control. Sections required for study in preparing for certification and licensing exams depend upon the license that an applicator is seeking. The categories, Ornamental & Turf, Pest Control Operator – Lawn & Ornamental (L&O), and Limited Lawn & Ornamental (Limited L&O) are responsible for material contained in the entire manual. Those seeking the Limited Commercial Landscape Maintenance certification are responsible for material presented in chapters 1 – 12 and chapters 22 – 24 only. 368 pp.

Weeds of Southern Turfgrasses - SP 79

Easy to use, practical weed identification guide contains 427 color photographs of 193 weed species with geographical range and life cycle descriptions. Included is a glossary of taxonomic terms. Indexed by common and scientific names.

Insects & Related Pests of Turfgrass in Florida - SP 140

Identify, learn about and control several insects and related arthropods that are common pests of turfgrass in Florida. Color photographs.

Troubleshooting Lawn Pests (Flashcard set) - SP 180

Learn to recognize organisms commonly found in Florida's turf. Forty-six laminated identification cards identify and describe insects and the damage they cause. Excellent field resource for turf and garden managers.

MCricket CD-ROM (SW-89)

The University of Florida knowledgebase on all ten species of mole crickets found in the United States, including Hawaii, Puerto Rico and the U.S. Virgin Islands. Covers life cycle, distribution, description, biological controls and damage. Includes a graphical identification key. Tutorials instruct the user in concepts of chemical and biological mole cricket control. Now on CD-ROM, plus many full-color photographs added. Runs on Macs and Windows-PCs with a CD-ROM drive and graphical World Wide Web browser software. Available on the WWW at > <http://www.ifas.ufl.edu/~ent1/mcricket/>

Pests In and Around the Home CD-ROM (SW-126)

The University of Florida's knowledgebase on pests of structures, lawns and landscapes. Contains information on biology, life cycle, identification, distribution, damage, management, and IPM. Contains links to hundreds of definitions, 150+ graphics and 300+ full-color photographs. Runs on Macs and Windows-PCs with a CD-ROM drive and graphical World Wide Web browser software.

Turfgrass Computer Training Tutorials (SW-121 and SW-127)

These two computer-verified training tutorials provide training on turfgrass pests. They are also authorized by the state of Florida for 1 CEU each for recertification purposes. Each contains 50 questions and provides the text and color photographs that the questions are based on. Requires Windows.

Applying Pesticides Correctly (SM 1)

This is the basic resource for the General Standards (Core) examination for pesticide applicators seeking to be certified and licensed to apply pesticides in Florida.

Call the **University of Florida Publications** office at 1-800-226-1764 during weekday office hours to place an order.

LICENSES FOR PERSONS WHO APPLY PESTICIDES TO TURF, LAWNS, AND ORNAMENTALS IN FLORIDA

Fred Fishel, Ph.D.
Pesticide Information Coordinator
IFAS, University of Florida

License Name	Pest Control Operator (PCO) - Lawn and Ornamental	Limited Commercial Landscape Maintenance	Limited Lawn and Ornamental	Ornamental and Turf
Statutory Authority	Chapter 482-Structural Pest Control Act	Chapter 482-Structural Pest Control Act	Chapter 482-Structural Pest Control Act	Chapter 487-Florida Pesticide Law
Responsible Agency and address.	FDACS, Bureau of Entomology & Pest Control 1203 Governor's Square Blvd., Suite 300 Tallahassee, FL 32301 850/617-7997	FDACS, Bureau of Entomology & Pest Control 1203 Governor's Square Blvd., Suite 300 Tallahassee, FL 32301 850/617-7997	FDACS, Bureau of Entomology & Pest Control 1203 Governor's Square Blvd., Suite 300 Tallahassee, FL 32301 850/617-7997	FDACS, Pesticide Certification Office Bureau of Compliance 3125 Conner Blvd., MD-1 Tallahassee, FL 32399-1650 850/617-7850
Who must have this license?	Businesses who perform pest control on lawns and ornamentals. Each business location must have a "certified operator-in-charge."	Commercial landscape maintenance personnel who apply certain pesticides to plant beds and ornamental plantings	<ul style="list-style-type: none"> • Government employees who apply pesticides to turf and ornamentals associated with government buildings. • Owners or employees of businesses who apply pesticides to the turf and ornamental plantings on their business property. 	Persons who apply or supervise the application of restricted use pesticides on golf courses, parks, cemeteries, and athletic fields.

License Name	Pest Control Operator (PCO) Lawn & Ornamental	Limited Commercial Landscape Maintenance	Limited Lawn & Ornamental	Ornamental & Turf
What kinds of pesticides trigger the licensing requirement?	License required for business to apply any pesticide including herbicides.	License required for application of pesticides and to perform integrated pest management on ornamental plants. Only those pesticides having the signal word, "Caution," on their labels may be applied. Insecticidal soaps, horticultural oils and Bt may also be applied.	License required for application of any pesticide including herbicides.	License required for application of restricted use pesticides, including herbicides.
Can the licensed applicator supervise unlicensed persons who work under his/her direct supervision?	The certified operator in charge may supervise an unlimited number of employees performing lawn & ornamental pest control from the business location. Each employee must have an identification card issued by FDACS-Bureau of Entomology & Pest Control.	Application by unlicensed persons not permitted. Each person who applies the pesticide must be licensed.	Application by unlicensed persons not permitted. Each person who applies the pesticide must be licensed.	The licensed applicator may supervise up to 15 unlicensed mixer/loaders and applicators at a time.
Qualification for license	The certified operator-in-charge must pass an examination.	<ul style="list-style-type: none"> • Applicator must pass an examination. 	<ul style="list-style-type: none"> • Applicator must pass an examination. 	<ul style="list-style-type: none"> • Applicator must pass an examination.
What are the qualifications to take exams?	<ul style="list-style-type: none"> • 3 years employment as a service employee of a licensed business that performs lawn and ornamental pest control, OR • a degree in entomology, horticulture, agronomy or related field PLUS 1 year experience working for a licensed firm OR • completion of a 1-year entomology program at a public university in FL which specializes in urban pest management and includes practical pest management experience. • a 2 year horticulture technology degree PLUS 1 year of employment as a service employee of licensed pest control business, OR • a specified number of credit hours in entomology, horticulture, etc. PLUS 1 year of employment as a service employee of a licensed pest control business. 	<ul style="list-style-type: none"> • Completion of 6 classroom hours of plant bed and ornamental continuing education training . 	<ul style="list-style-type: none"> • No qualifications 	<ul style="list-style-type: none"> • No qualifications

License Name	Pest Control Operator (PCO) Lawn & Ornamental	Limited Commercial Landscape Maintenance	Limited Lawn & Ornamental	Ornamental & Turf
License renewal	License renewed annually	License renewed annually	License renewed every 4 years	License renewed every 4 years
Recertification requirements	Annual recertification required. Recertify by: <ul style="list-style-type: none"> • Re-examination, OR • Obtain 2 core and 2 L&O classroom hours of acceptable continuing education units. 	Annual recertification required Recertify by: <ul style="list-style-type: none"> • Re-examination, OR • Obtain 4 classroom hours of acceptable continuing education units. 	Recertification required every 4 years. Recertify by: <ul style="list-style-type: none"> • Re-examination, OR • Obtain 4 classroom hours of acceptable continuing education units. 	Recertification required every 4 years. Recertify by: <ul style="list-style-type: none"> • Re-examination, OR • Obtain 4 core <i>and</i> 12 continuing education units by attending programs approved by Department
Fees	<ul style="list-style-type: none"> • \$300 testing fee • \$300 annual business license fee • \$150 annual certified operator-in-charge license fee 	<ul style="list-style-type: none"> • \$150 testing fee. Includes cost of 1-year license, if exam is passed. • \$75 annual license fee 	<ul style="list-style-type: none"> • \$150 testing fee. Includes cost of 4-year license, if exam is passed. • \$25 license renewal fee (4-years) 	<ul style="list-style-type: none"> • No testing fees. • \$100 for a 4-yr. Public license. • \$250 for a 4-yr. Commercial license
Insurance Requirements	Certificate of Insurance which meets requirements for minimum financial responsibility for bodily injury and property damage. <ul style="list-style-type: none"> • Bodily injury: \$100,000 each person & \$300,000 each occurrence; • Property damage: \$50,000 each occurrence & \$100,000 in the aggregate combined or single unit coverage; \$400,000 in the aggregate. 	Same as PCO	No insurance requirements.	No insurance requirements.

License Name	Pest Control Operator (PCO) Lawn & Ornamental	Limited Commercial Landscape Maintenance	Limited Lawn & Ornamental	Ornamental & Turf
What are the limitations on the license	Licensee cannot apply pesticides to golf courses, parks, cemeteries, or athletic fields.	Licensee cannot: <ul style="list-style-type: none"> • Operate a pest control business. • Apply pesticides to turf. • Apply pesticides to golf courses, parks, cemeteries, & athletic fields. Licensee is limited to: <ul style="list-style-type: none"> • Portable handheld 3-gallon compressed air or 5-gallon backpack sprayers. • Application of herbicides in plant beds and ornamental plantings and to IPM on ornamental plants using pesticides with caution label, insecticidal soaps, horticultural oils & B.T. 	Licensee cannot: <ul style="list-style-type: none"> • Operate a pest control business. • Apply pesticides to golf courses, parks, cemeteries, or athletic fields. 	Licensee is limited to pesticide applications to turf and ornamentals on golf courses, parks, cemeteries, & athletic fields.
Training requirements for persons working under direct supervision of the licensed applicator or certified operator	Each person under the direct supervision, direction, and control of certified operator must have at least 5 days of field training in lawn and ornamental pest control. In addition to this training each identification cardholder must receive 4 hours of classroom training in pesticide safety, integrated pest management, and applicable federal and state laws and rules with 6 months after issuance of the card or must have received such training within 2 years before issuance of the card. Each cardholder must receive at least 2 hours of continuing training in pesticide safety, integrated pest management, and applicable federal and state laws and rules by the renewal date of the card.	N/A. Each person making application must be licensed.	N/A. Each person making application must be licensed.	The licensed applicator must provide the following instruction and training to each unlicensed applicator working under their supervision: <ol style="list-style-type: none"> The safety procedures and precautions to be followed in using the product. The need to properly wear and maintain any required personal protective equipment. The common signs of pesticide poisoning. The dangers of eating, drinking or smoking while using pesticides. The need to wash clothing and bathe after working with pesticides. The name and location of a nearby medical facility that can provide emergency treatment for pesticide poisoning.. How and under what circumstances to immediately contact the licensed applicator under whose direct supervision the unlicensed person is working.

INSECT MANAGEMENT

Eileen A. Buss, Ph.D., Associate Professor and Landscape Entomologist

Several insects and mites feed on or live in grass, but not all of them cause economic or aesthetic damage. Many are harmless, some are beneficial, and some are pests. Some pests may need immediate control, especially if present in great numbers, but others may not be worth the time, effort, or cost of control. Feeding by pests may cause physical damage or just make the grass look temporarily bad. However, insects are only one of many potential causes for thin or brown grass. Diseases, nematodes, drought, nutritional disorders or incorrect chemical applications can also be damaging. Correct identification of the problem can save money and prevent unnecessary pesticide use. After the pest is correctly identified, information can be found on its life cycle, food preference, and habits. It is important to understand these things to properly time any corrective measures.

Scouting or monitoring for damage or pests is an important part of turfgrass management. Examine grass weekly in the spring, summer and fall, or train the mowing crew to record pest activity in areas that are often infested. Exactly how to monitor for each pest depends on where the insect lives or feeds.

Insects and their relatives can feed on the leaf tissue of grasses (e.g., various caterpillars), suck fluids from the leaves or crowns (e.g., southern chinch bugs, spittlebugs, scales, bermudagrass mites), and consume roots (e.g., white grubs, billbugs, mole crickets). In addition, mole crickets tunnel near the soil surface, uprooting grass plants and creating small mounds. Other nuisance organisms (e.g., ants, fleas, ticks, millipedes, chiggers, sowbugs) also occur in the turfgrass, but don't hurt the grass. Rather, they may bite, sting, or occasionally invade buildings.

Southern chinch bugs: To monitor, part the yellowing grass to look for moving insects on plants and in thatch. Vacuum declining areas with a Dustbuster or hand-held vacuum, and empty the filter to examine insects.

Mole crickets, caterpillars, scarab and billbug adults: Monitor by mixing 1 – 2 TBSP. (1.5 – 3 fl. oz.) of liquid dishwashing soap in 1 gallon of water; pour the solution onto 4 square feet near the damage. Insects will crawl to the surface if present in the grass, thatch, or upper soil layer. Examine several suspected areas. Mole cricket tunnels are also most visible early in the morning, when the dew is still present on the grass.

White grubs and billbug larvae: Watch for adult scarab beetles flying at night near lights from March to October. If a problem is suspected, cut 2-4 inches deep in a 1 foot square area of damaged grass. Lay the grass back, check root quality and look for grubs or billbug larvae in the soil. Many white grub species become damaging by late summer.

Cultural Controls: In general, healthy turf is less vulnerable to pests and can recover faster from an infestation. Follow the Best Management Practices for growing turfgrass. Mow at the correct height for the grass species, reduce thatch, and avoid over-watering. Avoid using flood lights or mercury vapor lights at night, especially in the spring when mole cricket adults, scarab beetles (adults of white grubs), and moths are flying because they are strongly attracted to light.

Natural Enemies: Various invertebrate predators (e.g., ground beetles, earwigs, spiders, and ants) and parasitoids (e.g., tiny wasps or flies) attack turfgrass pests. Although natural enemies rarely completely control pest populations, they do provide some natural suppression, so it is good to conserve them.

Insect Parasitic Nematodes: Insect parasitic nematodes in the genera *Steinernema* and *Heterorhabditis* help naturally suppress white grubs and mole crickets. Nematodes work better under moist soil conditions than in dry soils.

Insecticides: Most insecticides kill by either direct contact with the insect or by ingestion. Some may also exert a fumigating or vapor action under certain conditions. Products should be selected that will effectively control the pests without injuring the plants or result in another pest outbreak, and should minimize non-target impacts. Before using an insecticide, consider the following points:

- Select the right product. Only use an insecticide that is recommended to control the target pest and is labeled for the appropriate site (e.g., golf course, athletic field, residential area).
- Use the label rate or recommended amount. Too little won't control the pest; too much is illegal. **Read the container label carefully.**
- Apply it correctly. Thorough coverage is essential. The pesticide must reach the area of the plant where the pest is living and feeding. Many failures to control pests result from incorrect applications rather than product failure. Adding a wetting agent or spreader-sticker to a spray mixture may improve a pesticide's coverage and help provide greater control. When recommended (especially for white grubs), water the grass (¼ to ½ inch of water) immediately after treatment to move the insecticide into the root zone where the insects are feeding. Irrigating also brings insects closer to the soil surface, which increases their contact with the insecticide residues.

Insecticide Resistance Management: Resistance is reflected in the repeated failure of an active ingredient to effectively control the target pest when used according to the label recommendations. Product storage, application methods, and unusual climatic or environmental conditions should be ruled out first. Insecticide resistance management involves

monitoring pest population density, determining economic injury levels, and using integrated control strategies. If resistance is suspected, avoid retreating the population with a product from the same chemical class or mode of action group. Insecticides with different modes of action should be rotated.

Pest	Signs / Symptoms	Insecticides	Notes
Ants, Nuisance	Small mounds occur in turf, which may influence how a ball rolls on a golf course green. Ants may also invade buildings from the outside, or nest in trees or ornamental plant beds.	Abamectin Acephate ² Bifenthrin Chlorfenapyr Cyfluthrin Deltamethrin Hydramethylnon Lambda-cyhalothrin Permethrin Thiamethoxam	Ants that invade buildings may also have a nearby honeydew source – look for sap-feeding insects in nearby ornamental plants, shrubs, or trees. A combination of baits and broadcast applications may be necessary for control. To locate a hidden nest, leave some food out, watch where the ants trail, and aim the application near that trail or nest.
Ants, Imported Fire	These ants nest in large, sandy mounds that go deep into the soil. Any mound disturbance causes ants to immediately defend the nest. Stings result in tiny pustules, pain, itching, and sometimes an allergic reaction. See: http://fireant.tamu.edu/management/index.php	Abamectin Acephate ² Beta-cyfluthrin Bifenthrin Carbaryl ⁴ Chlorfenapyr Cypermethrin Deltamethrin Fenoxycarb Hydramethylnon Indoxacarb Lambda-cyhalothrin Permethrin Pyriproxifen Spinosad S-methoprene Sulfluramid Thiamethoxam	Scatter granules around the edge of the nest, not on top, for a mound treatment. Ants take the bait into the colony and feed the treated oils to each other, which results in colony death. Some baits work within 48 hours, some take a month. Fire ants that nest near buildings may be both an indoor and outdoor problem. Control is best achieved in the landscape before ants move indoors. Seal/caulk any cracks and crevices to prevent ant entry. Follow trails to find the nest; treat the area where ants are entering the building.
Beetles (adults)	Some adult beetles make small push-up mounds in turf and others just lay eggs in the plants or soil.	Acephate ² Bifenthrin Carbaryl ⁴ Cyfluthrin Deltamethrin Lambda-cyhalothrin	Adult beetles in the soil are not usually targeted for control, unless they are on the soil surface. Hatching larvae are more vulnerable to insecticides. Adult billbugs make small notches on grass stems. The hunting billbug is a major pest of bermudagrass and zoysiagrass. Adults are active at night and can be seen with a flashlight on the soil surface.

Pest	Signs / Symptoms	Insecticides	Notes
Billbugs (larvae)	<p>Young larvae feed in the stem and older larvae feed on roots. Patches of turf turn yellow, then brown. Grass stems break near the crown, and frass is visible in stems or stolons. Heavily infested sod falls apart when cut. Infestations may be misdiagnosed as early winter dormancy, slow spring green-up, or dollar spot disease.</p> <p>See EDIS publication ENY-321/LH037 http://edis.ifas.ufl.edu/lh037</p>	<p>Carbaryl⁴ Chlorantraniliprole Clothianidin Halofenozide Imidacloprid Thiamethoxam Trichlorfon</p>	<p>Billbug larvae are legless (white grubs have legs). They are present year-round in bermudagrass and zoysiagrass but can be hard to find. Overseeding with endophytic ryegrass reduces populations in the winter months. Monitor using the “tug test.”</p>
Caterpillars (armyworm, cutworm, grass loopers, tropical sod webworm)	<p>Young caterpillars skeletonize grass blades. Older caterpillars may notch the sides or completely eat the grass blades. Damaged grass may look ragged or scalped. In severe infestations, the ground may look like it is moving.</p> <p>Damage will be irregularly-shaped, unlike lawn mower/trimmer scalping.</p> <p>See EDIS publication ENY-352/IN608 http://edis.ifas.ufl.edu/in608</p>	<p>Acelepryn Acephate² <i>B. t.</i> var. kurstaki Bifenthrin Carbaryl⁴ Clothianidin Cyfluthrin Deltamethrin Diflubenzuron Dimilin Fipronil Halofenozide Indoxacarb Lambda-cyhalothrin Permethrin Spinosad Trichlorfon</p>	<p>Target young larvae with reduced-risk products like B.t., halofenozide, and spinosad. Caterpillars tend to become a problem in newly established turf, or in early fall, especially if the turf was fertilized heavily in late summer. Most feed at night. Turf can usually recover from caterpillar damage.</p>
Chiggers	<p>Chiggers are immature red mites that bite people, remain attached for 1 to 4 days, and cause severe itching. They tend to occur in areas of tight clothing.</p> <p>See EDIS publication ENY-212/IG085 http://edis.ifas.ufl.edu/ig085</p>	<p>Bifenthrin Carbaryl⁴ Cyfluthrin Deltamethrin Lambda-cyhalothrin Permethrin</p>	<p>Insecticidal control is difficult. Chiggers occur wherever their hosts live, including in lawn, picnic areas, and other recreational areas. Keep grass mowed and shrubs pruned.</p> <p>Prevent personal contact by using repellents, and wear protective clothing.</p>
Chinch Bugs, Southern	<p>Injured plants look stunted, yellowed, wilted, or dead. Small dead patches appear first, often near pavement or in stressed areas of St. Augustinegrass.</p> <p>See EDIS publication ENY-325/LH036 http://edis.ifas.ufl.edu/lh036</p>	<p>Bifenthrin Carbaryl⁴ Clothianidin Cypermethrin Deltamethrin Lambda-cyhalothrin Permethrin Thiamethoxam Trichlorfon</p>	<p>Many populations have become resistant to several insecticide chemical classes including pyrethroids. Use a high rate of insecticide with a wetting agent to penetrate thatch. Avoid using low rates in locations with reduced efficacy and combination products may also help reduce resistant populations. Rotate modes of action. Spot treat when possible. Reduce thatch thickness to minimize habitat.</p>

Pest	Signs / Symptoms	Insecticides	Notes
Fleas	<p>Fleas are external, blood-sucking parasites on the skin. The cat flea is most common.</p> <p>See EDIS publication ENY-205/IG087 http://edis.ifas.ufl.edu/ig087</p>	<p>Acephate Bifenthrin Carbaryl⁴ Cyfluthrin Cypermethrin Deltamethrin Esfenvalerate Lambda-cyhalothrin Permethrin</p>	<p>For best results, both the pet and lawn should be treated. Flea collars on pets may be necessary.</p> <p>Mow the lawn 1 or 2 days before treatment. Treat the entire area where pets normally sleep or play.</p> <p>Repeat as necessary.</p>
Greenbug (aphids)	<p>This aphid feeds on the phloem tissue of grasses and injects a toxin while feeding. The leaf area around the feeding site turns yellow and dies, sometimes turning burnt orange in color. Irregular dead patches may be surrounded by bands of yellow and rust-colored turf.</p> <p>http://entnemdept.ufl.edu/creatures/field/bugs/greenbug.htm</p>	<p>Acephate² Azadirachtin Bifenthrin Carbaryl⁴ Clothianidin Cyfluthrin Deltamethrin Dinotefuran Imidacloprid Lambda-cyhalothrin Permethrin Thiamethoxam</p>	<p>This is a major pest of sugarcane and wheat, but has been found on several warm season grasses. Populations can build rapidly. It has a history of insecticide resistance, so resistance management is important. Spot treat up to 3 ft around the visible injury, when possible.</p>
Ground Pearls	<p>These insects suck fluids from grass roots, which make irregular patches of turf look unhealthy. Grass yellows, browns, and dies, especially in hot, dry weather.</p> <p>See EDIS publication ENY-322/LH073 http://edis.ifas.ufl.edu/lh073</p> <p>See EDIS publication EENY-277/IN554 http://edis.ifas.ufl.edu/in554</p>	<p>None available</p>	<p>Ground pearls are often found by the nematode assay lab when they look for nematodes in soil samples. Properly fertilize, irrigate, and mow at the correct height for the turf species, to keep the turf growing ahead of the damage.</p>
Millipedes, Pillbugs, and Sowbugs	<p>These arthropods feed on decaying matter. They are occasional invaders in buildings but do not damage plants.</p> <p>See EDIS publication ENY-221/IG093 http://edis.ifas.ufl.edu/ig093</p>	<p>Bifenthrin Carbaryl⁴ Cyfluthrin Deltamethrin Lambda-cyhalothrin Permethrin</p>	<p>Widespread control is not usually recommended. Perimeter treatments may be needed, after gaps around doors and windows are sealed.</p> <p>Minimize mulch and moist areas.</p>

Pest	Signs / Symptoms	Insecticides	Notes
Mites	<p>Bermudagrass Mite – Infested turf has short leaves and internodes, resulting in a tufted or “witch’s broom” appearance. Grass leaf tips may be slightly yellowed. Large patches of turf may die, especially during hot, dry weather.</p> <p>For zoysiagrass mites, edges of grass blades look folded-over or curled. Spider mite-infested turf looks yellow or burned. Fine webbing and stippling damage may be present.</p> <p>See EDIS publication ENY-328/LH035 http://edis.ifas.ufl.edu/lh035</p>	<p>Bifenthrin Deltamethrin Lambda-cyhalothrin</p>	<p>Mow as low as possible, collect, and remove grass clippings to reduce the mite population. Using a wetting agent in the spray should improve coverage. Grass may outgrow damage if properly fertilized and irrigated. A repeat application may be necessary.</p>
Mole Crickets	<p>Most damage is caused by nymphs and adults tunneling in the soil, which exposes and dries out roots, and by root feeding. Tunnels are easiest to see in the morning, when dew is still on the grass. Damaged turf may thin, then die in large patches. Soil may feel spongy when walked on.</p> <p>Older nymphs make mounds later in the summer, when they come out at night and feed on grass blades.</p> <p>See EDIS publication ENY-324/LH039 http://edis.ifas.ufl.edu/lh039</p>	<p>Acephate² Bifenthrin Clothianidin Cyfluthrin Deltamethrin Fipronil Imidacloprid Indoxacarb Lambda-cyhalothrin Permethrin Thiamethoxam Trichlorfon</p> <p>Baits: Carbaryl⁴ Chlorpyrifos³ Indoxacarb</p>	<p>It is important to get insecticides into the soil, either by slit-injection, pre- or post-treatment irrigation (see product labels), or by using a wetting agent in the spray solution. Apply insecticides as late in the day as possible. Mole crickets are deeper in the soil during the day and closer to the soil surface at night. Use soap flushes to determine mole cricket age and density.</p> <p>Baits are most effective later in the summer, when older nymphs come onto the soil surface at night. Do not get baits wet.</p> <p>Beneficial nematodes attack large nymphs and adults, and do not damage plants. They are compatible with most insecticides, but not nematicides, to provide long-term mole cricket suppression.</p>
Scales and Mealybugs	<p>Scales and mealybugs may infest the leaves, crowns, or roots of turf plants. Leaves may first have spots or look yellowed, then turn brown, and die. Heavy infestations of rhodesgrass mealybug look like white fertilizer granules have caked around the grass nodes. Some sooty mold may be visible.</p>	<p>Bifenthrin Clothianidin Deltamethrin Distance IGR Imidacloprid Thiamethoxam</p>	<p>These insects are occasional turf pests, but they tend to be very damaging to groundcovers and ornamentals grasses. Cut infested leaf blades low and remove clippings to minimize infestations.</p>
Spittlebugs	<p>Damage on centipedegrass and St. Augustinegrass includes yellowing, purple streaking, browning, and turf death. Heavily infested turf feels “squishy” due to spittle masses in the thatch.</p> <p>See EDIS publication ENY-334/LH077 http://edis.ifas.ufl.edu/lh077</p>	<p>Bifenthrin Carbaryl⁴ Cyfluthrin Deltamethrin Lambda-cyhalothrin</p>	<p>Treat when most of the spittlebugs have become adults (June to September). Mow and irrigate before application to allow insects to penetrate the thatch.</p> <p>Spittlebugs cannot survive drought conditions. Avoid over-irrigation of turf to minimize infestation.</p>

Pest	Signs / Symptoms	Insecticides	Notes
Ticks	<p>Ticks are external parasites on skin. They can transmit diseases. Tiny seed ticks or engorged ticks may be seen attached to skin.</p> <p>See EDIS publication ENY-206/IG088 http://edis.ifas.ufl.edu/ig088</p>	<p>Bifenthrin Carbaryl⁴ Cyfluthrin Deltamethrin Lambda-cyhalothrin Permethrin</p>	<p>Insecticidal control is difficult. Keep grass mowed low to reduce humidity. Prevent personal contact with ticks by using repellents, wear protective clothing, and carefully inspect for and promptly remove any attached ticks.</p>
White Grubs	<p>White grubs live in the soil and feed on plant roots. Heavily infested turf may feel spongy when walked on, look yellowed or brown, and pull easily out of the soil. Sod may fall apart when cut. Animals may be seen feeding in an infested area. Swarms of parasitic wasps may hover just above infested turf. Identify grubs by the raster or hair patterns on the tip of their abdomens.</p> <p>See EDIS publication ENY-321/LH037 http://edis.ifas.ufl.edu/lh037</p>	<p>Preventative: Chlorantranilprole Clothianidin Dinotefuran Halofenozide Imidacloprid Thiamethoxam</p> <p>Curative: Carbaryl⁴ Trichlorfon</p> <p>Nematodes: <i>Heterorhabditis zealandica</i> <i>Steinernema glaseri</i></p>	<p>Apply preventative treatments when adult scarab beetles are laying eggs or when eggs start to hatch (April to June in most of Florida, for most species). Masked chafer damage appears in July. Sugarcane grub damage appears in September and October and may continue through January.</p>
Worms	<p>Worms may make small push-up mounds or castings in the turf.</p> <p>See EDIS publication CIR455/IN047 http://edis.ifas.ufl.edu/in047</p>	<p>None available</p>	<p>Control is not recommended. Worms help aerate the soil and are considered beneficial organisms.</p>

TURFGRASS INSECT AND NUISANCE PEST CONTROL NOTES

¹Only a few formulations of recommended insecticides are listed to serve as examples. Many others are available. No endorsement of products is intended, nor is criticism of unnamed products implied. **Read container label carefully for use directions, application techniques, irrigation requirements, worker protection information, and precautions. Be sure** the formulation of pesticide you use is labeled for use on turfgrass.

²When using acephate, check pH of spray water and adjust to 5.5 - 6.0 when pH is above 7.0. Acephate is not registered for use on residential turf except as a fire ant mound treatment. Acephate will still be registered for broadcast application to turf on golf courses and sod farms.

³Dursban not labeled for residential use.

⁴Sevin (carbaryl) insecticide is going through re-registration with the EPA. Until this process is completed, all home lawn uses had to be removed from the labels of liquid Sevin products. Commercial lawn and ornamental uses are still labeled. New product labeling is expected within the next 6 months.

Common insecticides listed by chemical classes for turf and/or ornamental use in Florida.

IRAC Mode of Action Classification	Chemical Classes	Mode of Action	Active Ingredients / Chemical Names / Trade Name Examples¹
1A	Carbamates	Acetylcholine esterase inhibitor	Carbaryl (Sevin)
1B	Organophosphates	Acetylcholine esterase inhibitor	Acephate (Orthene), chlorpyrifos (Dursban), diazinon, dimethoate (Cygon), malathion, trichlorfon (Dylox)
2A	Cyclodiene organochlorines	GABA-gated chloride channel antagonists	Chlordane, dicofol (Kelthane), lindane
2B	Phenylpyrazoles	GABA-gated chloride channel antagonists	Fipronil (Chipco Choice, Chipco TopChoice, MaxForce FC)
3	DDT, pyrethroids, pyrethrins	Sodium channel modulators	Bifenthrin (Talstar, Onyx), beta-cyfluthrin, cyfluthrin (Tempo), cypermethrin (Demon), deltamethrin (DeltaGard), cyhalothrin, lambda-cyhalothrin (Scimitar), esfenvalerate, fenpropathrin, fenvalerate, permethrin (Astro), resmethrin
4A	Neonicotinoids	Nicotinic acetylcholine receptor agonists / antagonists	Acetamiprid (TriStar), clothianidin (Arena), dinotefuran (Safari), imidacloprid (Merit), thiamethoxam (Meridian)
5	Spinosyns	Nicotinic acetylcholine receptor agonists (allosteric) - not group 4	Spinosad (Conserve)
6	Avermectins	Chloride channel activators	Abamectin (Ascend, Clinch, Varsity Fire Ant Bait)
7A	Juvenile hormone analogs	Juvenile hormone mimics	Hydroprene, kinoprene, methoprene (Extinguish)
7B	Fenoxycarb	Juvenile hormone mimics	Fenoxycarb (Award Fire Ant Bait)
7C	Pyriproxyfen	Juvenile hormone mimics	Pyriproxyfen (Distance Fire Ant Bait, Distance IGR, Esteem)
8A	Alkyl halides	Compounds of unknown or non-specific mode of action (fumigants)	Methyl bromide
11B2	<i>B. t. subspecies kurstaki</i>	Microbial disruptors of insect gut membranes	<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> (Dipel)
15	Insect growth regulator	Inhibitor of chitin biosynthesis	Diflubenzuron (Dimilin)
18A	Diacylhydrazines	Ecdysone agonists / molting disruptors	Halofenozide (Mach 2), tebufenozide (Confirm)
18B	Azadirachtin	Ecdysone agonists / molting disruptors	Azadirachtin (Azatrol, Azatin)
20A	Hydramethylnon	Mitochondrial complex III electron transport inhibitors (Coupling site II)	Hydramethylnon (Amdro)
22	Oxadiazine	Voltage-dependent sodium channel blockers	Indoxacarb (Advion, Provaunt)
28	Anthranilic diamide	Acelepryn depletes calcium from insect muscles disrupting normal contraction.	Chlorantraniliprole (Acelepryn, Calteryx)

¹Specific products are listed for example only. Neither inclusion of products nor omission of similar alternative products in this publication is meant to imply any endorsement or criticism.

DISEASE MANAGEMENT

Philip F. Harmon, Ph.D.

Associate Professor and Extension Plant Pathologist

Diseases can disfigure turfgrass around homes, recreational areas, and commercial grounds. Fortunately, grasses receiving proper cultural practices including proper irrigation, mowing, and fertilizing are less likely to develop diseases and are not as likely to be seriously damaged if a disease occurs. By enhancing plant vigor, diseases will be minimized and the need for the use of costly fungicides will be reduced. Submit suspected disease samples to a University of Florida Extension Plant Disease Clinic for diagnostic confirmation of disease problems and to receive management recommendations. If used, alternate between classes of fungicides to prevent development of fungicide-resistant pathogens. See the end of this section for a table that lists fungicide chemical classes. Read labels carefully. Many fungicides are limited regarding the turfgrass site on which they may be used. Also, many fungicides have a limit on the amount and/or number of applications allowed within a one-year period.

Disease	Affected Grasses	Symptoms	Cultural Controls	Fungicides ¹
Brown Patch Large Patch Rhizoctonia Blight <i>(Rhizoctonia solani)</i>	Bahiagrass Bermudagrass Carpetgrass Centipede Ryegrass Seashore paspalum St. Augustinegrass Zoysiagrass	Grass is killed in circular to irregular areas that may expand to several feet in diameter. Leaf fascicles pull easily from plant due to rot at leaf base. Occurs during humid, rainy weather. High N, thatch buildup, and excessive moisture favor disease. St. Augustine, carpet, centipede, zoysia (fall through spring), and rye-grasses (winter) are more affected. This is usually not a summer disease.	Maintain adequate fertility. Avoid excess fast-release nitrogen. Irrigate deeply. Reduce thatch.	azoxystrobin chlorothalonil ⁴ fenarimol ² fludioxonil fluoxastrobin flutolanil iprodione ⁴ Junction ³ mancozeb ⁴ metconazole ² myclobutanil ² polyxin D propiconazole ² pyraclostrobin thiophanate-methyl thiram ⁵ triadimefon ² trifloxystrobin triticonazole ² vinclozolin
Rhizoctonia Leaf and Sheath Spot <i>R. oryzae, R. zeae</i>	Bermudagrass Seashore paspalum	Occurs during summer months when weather is hot and humid. The most commonly observed symptoms are necrotic rings or partial rings that vary from a few inches to a few feet in diameter. Basal leaf area is <u>not</u> rotted. Spots may be observed on leaves at edge of rings.	Unknown at this time, but increasing nitrogen level may be useful.	azoxystrobin chlorothalonil ⁴ fludioxonil fluoxastrobin flutolanil iprodione ⁴ mancozeb ⁴ pyraclostrobin thiram ⁴ The above products may be useful for control.

Disease	Affected Grasses	Symptoms	Cultural Controls	Fungicides ¹
Cercospora Leaf Spot (<i>Cercospora fusimaculans</i>)	St. Augustinegrass	Brown to purple leaf spots in patches 2-3" in diameter. In high disease severity, entire leaves will yellow, wither and die. Warm, humid weather favors disease incidence.	N may reduce disease. Water deeply only when needed in mornings. 'Bitter-blue' selections are more resistant.	None available. Fungicides used to control other leaf spot diseases will provide suppression.
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	Bahiagrass Bermudagrass Centipedegrass Ryegrass Seashore paspalum St. Augustinegrass Zoysiagrass	On fine textured grasses, spots appear 1-2" in diameter. Spots larger on coarse grasses. Leaves develop marginal, irregular, light tan lesions with reddish brown borders. Active during 60-80°F in fall through spring. Moisture from fog, dew, or irrigation initiate disease. Low soil moisture, thatch, low N and K favor disease.	Avoid N deficiency. Irrigate in morning. Avoid thatch buildup.	boscalid chlorothalonil ⁴ fenarimol ² fluoxtrobin iprodione ⁴ Junction ³ mancozeb ⁴ metconazole ² myclobutanil propiconazole ² pyraclostrobin thiophanate-methyl thiram ⁴ triadimefon ² triticonazole ² vinclozolin
Fairy Ring (<i>Chlorophyllum</i> sp., <i>Marasmius</i> sp., and others) <i>Also see localized dry spots/rings</i>	All grasses	<u>Type I:</u> Dead rings (see Localized Dry Spots / Rings). <u>Type II:</u> Irregularly sized circular to semi-circular bands of lush green turf become apparent. Turf within circular area may decline, brown and thin. Mushrooms may be present. <u>Type III:</u> Mushrooms present, but grass is unaffected.	Difficult to control. Plugging or aerating to allow more water and fertilizer to reach the roots may help. Additional fertilizer will mask Type II rings.	azoxystrobin fluoxastrobin flutolanil metconazole ² pyraclostrobin triadimefon triticonazole ² *If mushrooms are present, collect and discard into garbage.

Disease	Affected Grasses	Symptoms	Cultural Controls	Fungicides ¹
Gray Leaf Spot (<i>Pyricularia grisea</i>)	Centipedegrass St. Augustinegrass	Small brown to ash-colored leaf spots with purple to brown margins. Lesions become covered with the gray, velvety, fungal mycelium of <i>Pyricularia grisea</i> . In severe cases leaves appear scorched. Prevalent during rainy, summer months. Mainly on St. Augustinegrass.	Avoid excess N. Irrigate deeply in early morning. Reduce traffic. Mostly a problem on recently planted or atrazine-treated stressed St. Augustinegrass.	azoxystrobin chlorothalonil ⁴ fluoxastrobin mancozeb ⁴ metconazole ² polyoxin D (<i>suppression only</i>) propiconazole ² pyraclostrobin thiophanate-methyl triadimefon ² trifloxystrobin
“Helminthosporium Leaf Spot” (<i>Bipolaris</i> , <i>Drechslera</i> , <i>Exserohilum</i> spp.)	Bahigrass Bermudagrass Ryegrass Seashore paspalum St. Augustinegrass Zoysiagrass	Symptoms include a leaf spot and 'melting-out' phase. Small oblong purplish to brown leaf spots. Numerous lesions cause leaves to turn reddish-brown and die. Sheath and crown rot may be present. Rye and bermuda are most susceptible. Most prevalent when temperatures range from 68-95°F during mild periods of fall through spring.	Maintain a balanced fertility. Irrigate deeply in the mornings. Raise mower height during disease outbreaks. Reduce thatch. Increase K in areas where disease is known to occur.	azoxystrobin chlorothalonil ⁴ fludioxonil fluoxastrobin iprodione ⁴ mancozeb ⁴ myclobutanil propiconazole ² pyraclostrobin trifloxystrobin triticonazole ² vinclozolin
Pythium Blight (<i>Pythium</i> spp.)	All grasses used for overseeding.	Grass dies in spots or streaks. Initially, the affected grass has a dark color and a greasy appearance. After prolonged moist or foggy periods, the cottony mycelium may be seen on the turf. <i>Pythium</i> can be spread by foot traffic or mowers passing over infected grasses. Occurs during warm, humid, weather after the grass is established.	Improve aeration and drainage. Avoid frequent, shallow irrigation. Reduce mowings and minimize equipment or foot traffic across infected turf. Wash equipment that passes from infected to non-infected grass areas.	azoxystrobin chloroneb cyazofamid etridiazole fluopicolide fluoxastrobin fosetyl-Al Junction ³ mancozeb ⁴ mefenoxam phosphorous acid propamocarb hydrochloride pyraclostrobin <i>To minimize the potential for resistance, alternate between classes of fungicides.</i>

Disease	Affected Grasses	Symptoms	Cultural Controls	Fungicides ¹
Pythium Damping-off <i>(Pythium spp.)</i>	All grasses used for overseeding.	Seed fails to germinate or germination is erratic. Seedlings killed after emergence have water-soaked lesion at the soil surface.	Provide good seed-soil contact to ensure rapid germination. Monitor water closely.	Apron XL (Seed treatment) Fungicides listed for Pythium Blight are useful, but should be applied <u>after</u> seeding.
Pythium Root Rot <i>(Pythium spp.)</i>	All grasses	Roots are dark, soft with few or no feeder roots present. Root rot is favored in poorly drained or continuously wet soils. Areas will appear chlorotic and be less vigorous in growth, but usually do not die. Can occur year around, especially on over-irrigated sites.	Avoid overwatering. Aerate compacted and poorly drained soils. Foliar fertilizer treatments may be useful.	azoxystrobin chloroneb cyazofamid etridiazole fluopicolide fluoxastrobin fosetyl-Al phosphorous acid propamocarb hydrochloride pyraclostrobin <i>To minimize the potential for resistance, alternate between classes of fungicides.</i> Except for fosetyl-Al, water into the root-zone.
Rust <i>(Puccinia spp.)</i>	St. Augustinegrass Zoysiagrasses	Small yellow to orange or reddish-brown pustules on the leaves during mild, humid weather. Heavily infected area appears thin and chlorotic. Rye- and zoysia-grasses are most susceptible.	Plant resistant or tolerant varieties. Maintain rapid growth by fertilizing. Mow frequently and remove clippings.	azoxystrobin fluoxastrobin Junction ³ mancozeb ⁴ metconazole ² myclobutanil triadimefon ² trifloxystrobin triticonazole ² propiconazole ² pyraclostrobin
Slime Mold <i>(Physarum sp., and Fuligo sp.)</i>	All grasses	Grass is covered with gray to black soot-like growth or prominent white or yellow masses during warm, moist weather. Slime molds do not injure turf.	Brush off or wash off the mold with a strong stream of water. Mow.	mancozeb

Disease	Affected Grasses	Symptoms	Cultural Controls	Fungicides ¹
<p>Bermudagrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)</p> <p>Take-all Root Rot (same pathogen as above)</p>	<p>Bermudagrass</p> <p>Seashore paspalum St. Augustinegrass</p>	<p>Disorder first appears as chlorotic patches 8-24" in diameter. Without control, patches will expand. Grass thins and develops a bare spot. Green shoots next to chlorotic ones are common. Plants in the affected areas have poor root system, no rhizomes and very few stolons. Usually observed first on outside edge of golf course putting greens. Primarily observed in summer and fall.</p>	<p>Raise mower height by 50% to increase photosynthetic area. Do not scalp St. Augustinegrass when mowed. Increased fertility may help by encouraging rapid cover of affected areas. Foliar fertilizer applications may be useful. Topdress golf course greens frequently. Alleviate all stresses on the grass.</p>	<p>Some <u>preventive</u> control of 'patch' and 'decline' type diseases has been achieved by use of products containing azoxystrobin, myclobutanil, propiconazole, thiophanate-methyl, triadimefon², fenarimol, fluoxastrobin, and pyraclostrobin. Use only preventive rates of triadimefon², propiconazole, and myclobutanil on Bermudagrass. The DMI fungicides are likely to have a negative impact on bermudagrass putting greens when used more than once. Preventive means at least one month <i>prior</i> to development of disease symptoms. Propiconazole would not be recommended on putting greens for the summer months (see footnote #2). Water into the root zone.</p>
<p>Anthracnose (<i>Colletotrichum graminicola</i>)</p>	<p>All grasses</p>	<p>The causal fungus can infect leaves, sheaths, and tillers. Leaf infection appears as reddish-brown to brown lesions that are often surrounded by a yellow halo. Lesion size may span the blade width and often one lesion will cause complete yellowing of a blade. Tiller infection results in stem girdling and the subsequent appearance of small, yellow patches of turf. The causal fungus can be observed with a hand lens. It will appear as a dark, cushion-like reproductive structure (acervulus) with black spines (setae) extending from the cushion.</p>	<p>Avoid stressed turf caused by pests, fertility imbalances, or moisture extremes. Thatch removal will be helpful.</p>	<p>azoxystrobin fluoxastrobin chlorothalonil⁴ fenarimol² fludioxonil metconazole² myclobutanil² propiconazole² pyraclostrobin thiophanate-methyl triadimefon² trifloxystrobin triticonazole²</p>
<p>Localized Dry Spots/Areas/Rings (Basidiomycete fungi, primarily <i>Lycoperdon</i> spp.)</p>	<p>Bermudagrass putting greens, especially those less than 4 yrs. old</p>		<p>The water-repelling (hydrophobic) soil must be broken up and wetted with irrigation and soil wetting agents. When fairy ring fungi are involved, applications of fungicides and wetting agents will help alleviate symptoms.</p>	<p>Soil wetting agents</p> <p>azoxystrobin fluoxastrobin flutolanil pyraclostrobin</p>

Disease	Affected Grasses	Symptoms	Cultural Controls	Fungicides ¹
Algae (various species; primarily blue-green species on surface)	All grasses Most prevalent on putting greens & other turf mowed too short.	Turf areas in partially shaded, damp locations become weak and begin to thin. Long-term overcast, rainy weather periods encourage algae on putting greens. These algae are commonly green or brown in color and can be sheet-like, leaf-like, or cushion-like in appearance. Due to their high water content, algae are often quite slippery. Algae growth may become so prolific that they cover turf plants and inhibit irrigation penetration.	Improve air circulation and light exposure. Improve drainage and reduce irrigation frequency and amount. Reduce freely available nitrogen at site. On putting greens, verticut or aerify to disrupt algal mats. Topdress frequently.	chlorothalonil ⁴ Junction ³ mancozeb ⁴

DISEASE CONTROL NOTES

¹Only single active ingredient products are listed. Many companies have products that are mixtures of two active ingredients. Presence of a fungicide in this list does not constitute a recommendation. Trade names are used with the understanding that no endorsement is intended nor is criticism implied of similar products which are not mentioned. All chemicals should be used in accordance with the manufacturer's instructions. Do not add adjuvants, surfactants, etc. to fungicides unless specified by the label.

²Bermudagrass may exhibit phytotoxicity to propiconazole and other DMI fungicides. See labels.

³Phytotoxicity may occur depending on turfgrass varietal differences and with multiple applications of Junction. Apply recommended rate to small area and observe for 7 to 10 days for signs of injury. If phytotoxicity occurs, discontinue use.

⁴Chlorothalonil, iprodione, thiram, mancozeb, chloroneb, and vinclozolin cannot be used on residential (home) lawns. Some can be used on sod, business, industrial, and golf course turfgrass sites. See label for applicable restrictions.

Turfgrass fungicides listed by chemical class for use in Florida.

Chemical Group	Common Name¹ (Trade Name Example²)	Location of Activity	Mode of Action	Mode of Action FRAC Codes³
Acylalanines (PhenylAmides)	Mefenoxam (Subdue)	Systemic; upward movement	Nucleic acid synthesis	4
Acylpicolide	Fluopicolide (Stellar)	Contact; Systemic; upward movement	Delocalisation of spectrin like proteins	43
Aromatic Hydrocarbons	Chloroneb (Tersan) Etridiazole (=Ethazole) (Terrazole)	Contact	Lipids and membrane synthesis	14
Carbamates	Propamocarb (Banol)	Systemic; upward movement	Lipids and membrane synthesis	28
Carboxamides	Boscalid (Emerald) Flutolanil (ProStar)	Systemic; upward movement	Respiration (complex II)	7
Chloronitriles	Chlorothalonil (Daconil)	Contact	Multiple sites	M5
DeMethylation Inhibitors	Fenarimol (Rubigan) Metconazole (Tourney) Myclobutanil (Eagle) Propiconazole (BannerMaxx) Triadimefon (Bayleton) Triticonazole (Trinity, Triton)	Systemic; upward movement	Sterol biosynthesis in membranes	3
Dicarboximides	Iprodione (Chipco 26GT, Iprodione Pro) Vinclozolin (Curalan)	Local-penetrant	Lipids and membrane synthesis	2
Dithiocarbamates	Mancozeb (Dithane, Fore) Thiram	Contact	Multi-site contact activity	M3
Inorganic Metals	Copper Hydroxide	Contact	Multi-site contact activity	
Phosphonates	Fosetyl-Al (Alette, Chipco Signature) Phosphorous Acid (Alude, Resyst, Magellan, Vital)	Systemic	Unknown	33
Polyoxins	Polyoxin D zinc salt (Endorse)	Systemic; upward movement	Glucan and cell wall synthesis	19
PhenylPyrroles	Fludioxonil (Medallion)	Contact	Signal transduction	12

Chemical Group	Common Name ¹ (Trade Name Example ²)	Location of Activity	Mode of Action	Mode of Action FRAC Codes ³
QoI quinone outside inhibitors	Azoxystrobin (Heritage) fluoxastrobin (Disarm) Pyraclostrobin (Insignia) Trifloxystrobin (Compass)	Systemic; upward movement Mesostemic	Respiration (complex III)	11
QiI-quinone inside inhibitor	Cyazofamid (Segway)			
Thiophanates (MBC fungicides)	Thiophanate methyl (3336)	Systemic; upward movement	Mitosis and cell division	1

¹Read all labels to determine the location where it is legal to use the products on turfgrass. For example, some products can only be used on golf courses, whereas others can be used on all turf sites except residential turfgrass.

²Specific products are listed for example only. Neither inclusion of products nor omission of similar alternative products in this publication is meant to imply any endorsement or criticism.

³FRAC = Fungicide Resistance Action Committee. Codes indicate the biochemical target site. M3, M4, and M5 indicate multisite inhibitor (broad mode of action) with no significant risk of resistance. See www.frac.info for further information. When considering rotation and tank mixes, be sure to use materials that do not have the same mode of action.

NEMATODE MANAGEMENT

William T. Crow, Ph.D., Associate Professor and Landscape Nematologist

Nematodes are important pests that commonly contribute heavily to the decline of turf in Florida. However, many times weak turf growth is blamed on nematodes when poor cultural practices, fungi, insects, nutrient problems, soil compaction, poor drainage, or other environmental problems may be the actual cause. Correct diagnosis is important before using of a nematicide. Nematicides vary in their effectiveness against different species of nematodes. Also, no nematicide can be used on all kinds of sites; consult label carefully to be sure a product can be used on a particular site.

DIAGNOSIS

ABOVE-GROUND SYMPTOMS: wilting and slow recovery from wilt; yellowing; decline or “melting out;” irregular shaped areas of declining turf; weed invasion.

ROOT SYMPTOMS: short roots with few branch roots compared to healthy roots; dark color, sometimes with swollen root tips; reduced root system that does not hold soil together when plugs or cores are lifted from the sod.

NEMATODE SAMPLING is the key to correct diagnosis. Instructions and forms for sample submission to the University of Florida Nematode Assay Lab can be obtained at: <http://nematology.ifas.ufl.edu/assaylab>.

NEMATODE MANAGEMENT TACTICS

IMPROVE TURF MANAGEMENT PRACTICES. If nematode population is high, address the problem with an integrated program of improved cultural practices, planting a different kind of grass if that is warranted and feasible, and chemical control where it is legal and practical. Most grasses can withstand moderate numbers of most kinds of nematodes. Deep, infrequent watering encourages deeper rooting of the turf, enabling grass to reach more water and nutrients than turf having a short root system due to frequent shallow watering. However, once nematode damage occurs increased irrigation may be required to prevent turf decline. Avoid excess nitrogen fertilization, as this encourages lush, succulent roots conducive to nematode population increases. Instead, use the recommended amounts of fertilizer, but split the amount into more frequent applications. Avoid unnecessary stresses to turf such as mowing too short. Alleviate compacted soils, poor drainage, and other soil physical problems, and correct any nutrient deficiencies.

PLANT A DIFFERENT GRASS. Planting another type of grass (see Table 1) may be a practical choice if the new grass provides acceptable quality, but no variety of any turfgrass is known to have true resistance to all nematodes. Using proper turf management practices (see above) is a more practical approach.

CHEMICAL NEMATICIDES can sometimes give turf short-term relief from stress caused by nematodes. Some are very toxic to plants as well as nematodes and other animal life, so must be used to treat soil to reduce nematode populations and other soil-borne pests before planting. Others are relatively safe for living turf, and can be applied to established grass to reduce nematode activity while the grass is growing. All nematicides are relatively toxic to people, pets, and wildlife, and all are quite soluble in water, so they pose serious threats to people and the environment if used carelessly. Therefore, they are Restricted Use Pesticides for most situations, and their use is strictly regulated.

EFFECTS OF NEMATICIDES ARE ONLY TEMPORARY. Fumigants leave behind no residual active ingredients, so nematodes that survived the treatment (e.g., were too deep to be reached by it or were protected inside fresh roots) or were brought in on new sod or sprigs can begin to re-colonize the normal turf root-zone immediately. The non-fumigant nematicides that may be applied to living turf must remain in the root-zone (top 3-4) inches in which most turfgrass roots grow) for several weeks to be effective. However, they will eventually dissipate from that region as a result of combined effects of leaching and decomposition. Once the chemical is gone, there are usually some nematodes ready to resume feeding and reproducing. With either kind of nematicide, the treatment only provides a limited period of relief from nematode stress. The treatment cannot result in the desired improvement of turf health unless other stresses are also controlled and the nutrients and water that are needed for good root growth are available.

Table 1. Nematodes and the grasses most affected by each.

Turfgrass	Sting ¹	Lance ²	Stubby-root ³	Spiral ⁴	Ring ⁵	Root-knot ⁶	Cyst ⁷
Bahiagrass	X	X					
Centipedegrass	X		X		X		
St. Augustinegrass	X	X	X			X	X
Bermudagrass	X	X	X			X	
Zoysiagrass	X	X	X			X	
Seashore paspalum	X	X	X	X		X	

¹Sting nematodes damage all grasses; generally found only in very sandy soils.

²Lance nematodes are widely distributed and attack all turfgrasses in Florida.

³There are two types of stubby-root nematodes commonly associated with turfgrasses in Florida, *Paratrichodorus minor* is more common on bermudagrass whereas *Trichodorus proximus* is more common on St. Augustinegrass. St. Augustinegrass appears to be more damaged by stubby-root nematodes than bermudagrass.

⁴Several genera of spiral nematodes are found frequently with turfgrasses. Some genera are considered damaging, while others are not.

⁵Ring nematodes are widely distributed and found associated with all turfgrass types, but are considered important pests only on centipedegrass.

⁶Root-knot nematodes found frequently on most turf types. Their effects on Florida turf are not well known, but they are believed to be injurious at high population densities.

⁷Cyst nematodes normally attack only St. Augustinegrass, and are found most commonly on the east coast and central Florida; high populations can damage this grass severely, and cyst nematodes are very hard to control with chemicals.

SOIL FUMIGATION BEFORE PLANTING. Multi-purpose soil fumigants (Table 2) can be used to treat planting sites before planting new turf or during renovation, to promote rapid and uniform establishment of new turf. Fumigants reduce numbers of nematodes and some soil-borne fungi, insects, and weeds. A loose, open-pored soil permits rapid and uniform diffusion of fumigant vapors. Moderate soil moisture is best: water-filled pores inhibit diffusion of the gas, while very dry soils allow fumes to escape too quickly, which also renders the application ineffective. Soil temperatures should be in the 50°-80°F range.

NEMATICIDES FOR ESTABLISHED TURFGRASS.

Nortica: The bionematicide Nortica contains a bacterium that colonizes turf roots and protects them from nematode damage for a period of time. UF research has found consistent improvement in turf root growth following treatment with Nortica prior to the onset of summer stress.

1,3-Dichloropropene: Curfew Soil Fumigant is a fumigant nematicide that is slit-injected into established turf on golf courses and athletic fields. It is very effective against sting nematode and other nematodes in the soil. It is not systemic, so generally will not kill nematodes inside of turf roots. Curfew is only applied by custom applicators. Telone II has the same active ingredient as Curfew and can be used on sod farms. Telone II should not be applied to mature sod because the injection process slices the sod and can affect harvestability.

Multiguard Protect: This nematicide/fungicide is labeled for use on golf course tees, greens, and fairways and sod farms. This is a new class of nematicide having furfural as the active ingredient. After the initial irrigation to water the product in, do not irrigate again for at least 24 hours. Usually several applications (3 to 4 wk apart) are required to achieve good nematode reduction.

MustGro Invest: This new bionematicide will be launched sometime in 2012. This is a granular product made of formulated mustard plant material. When it gets wet it releases a nematicidal gas (allyl-isothiocyanate) that is moved into the soil with irrigation. The turf surface should be dry (no dew or moisture) when MustGro Invest is applied, and then the turf should be irrigated immediately following application. Based on UF research, the product will work best when used on areas that have an efficient irrigation system and good soil infiltration. Optimally, MustGro Invest should be applied immediately following aeration. MustGro Invest contains 5% nitrogen, so it has fertility as well as nematode effects.

Other Nematode Management Products: There are several other nematode management products available, most are microbially or botanically derived. While certain of these may help in some instances, none are currently recommended by the University of Florida because scientific evidence of their efficacy against nematodes in the field is either minimal or non-existent. The University of Florida is continuing to work with some of these in order to gather research data, so some may be recommended in the future.

Table 2. Multi-purpose soil fumigants for treating turf planting sites before planting. All are Restricted Use Pesticides for most applications.

Fumigant Products	Comments
Telone C-17	This liquid fumigant is injected into the soil with tractor-mounted equipment. Maximum effectiveness is achieved when soil is covered with a plastic tarp for one to several days.
Metham-sodium products	Apply these water-soluble materials either as a drench or spray in water or inject through tractor-mounted chisels. Cover with a plastic tarp after application, for maximum benefit.
Methyl bromide / chloropicrin mixture (many brands)	Inject through chisels and cover immediately with a plastic tarp for safety and effectiveness. Usually done by custom applicators who have the special equipment and training to handle these especially toxic pesticides.

Table 3. Risk thresholds for most warm-season turfgrasses used by the University of Florida Nematode Assay Laboratory

“-” = not believed to cause significant damage.

“M” = Turf is considered at moderate risk of damage. Damage may become evident if the turf is placed under stress conditions.

“H” = Turf is considered at high risk of damage. Root systems are likely damaged and turf quality may be declining.

Turfgrass Species	Root-knot (<i>Meloidogyne</i>)		Sting (<i>Belonolaimus</i>)		Lance (<i>Hoplolaimus</i>)		Subby-root (<i>Paratrichodorus</i>)		Subby-root (<i>Trichodorus</i>)		Spiral (<i>Helicotylenchus</i>)		Spiral (<i>Peltamigratus</i>)		Ring (<i>Mesocriconema</i>)		Sheath (<i>Hemicyclophora</i>)		Sheathoid (<i>Hemicriconemoides</i>)		Awl (<i>Dolichodorus</i>)		Cyst (<i>Heterodera</i>)	
	M	H	M	H	M	H	M	H	M	H	M	H	M	H	M	H	M	H	M	H	M	H	M	H
Bermudagrass	80	300	10	25	40	120	150	300	40	120	700	1500	150	300	500	1000	150	300	500	1000	10	25	-	-
Ultradwarf Bermudagrass	40	200	5	20	40	120	150	300	40	120	700	1500	150	300	500	1000	150	300	500	1000	5	20	-	-
Zoysiagrass	80	300	10	25	40	120	150	300	40	120	700	1500	150	300	500	1000	150	300	500	1000	10	25	-	-
Seashore paspalum	80	300	10	25	40	120	150	300	40	120	300	700	150	300	500	1000	150	300	500	1000	10	25	-	-
St. Augustinegrass	80	300	25	50	40	120	40	120	40	120	700	1500	150	300	500	1000	150	300	500	1000	10	25	10	40
Centipedegrass	80	300	10	25	40	120	150	300	40	120	700	1500	150	300	150	300	150	300	150	300	10	25	-	-

*These nematodes thresholds are based upon numbers per 100 cc of soil extracted using a sugar-flotation with centrifugation method.

**While bahiagrass is a host for many of these nematodes, it is very tolerant to them and seldom is damaged. Therefore, no thresholds are given.

***Other nematodes such as dagger, lesion, stunt, etc. may damage turf in Florida, but damage from these is very rare so thresholds are not listed.

****These thresholds are based upon nematodes, grasses, and conditions in Florida only. They may not apply in other states.

WEED MANAGEMENT

J. Bryan Unruh, Ph.D. and Barry J. Brecke, Ph.D.
Extension Turf Specialist and Research Weed Scientist

The best defense against weeds is a dense, vigorously growing turf. By adapting the right grass to the site and following correct cultural management, including proper fertilization, mowing, and irrigation, weeds will not be able to compete as well as with the turf. Before deciding to use any weed control, diagnose first why the turf is thin and weeds are invading. Correct the basic problem of unhealthy turf before using any weed control. **HERBICIDES ARE NOT A SUBSTITUTE FOR SOUND CULTURAL PRACTICES.**

The first step toward a successful weed management program is the accurate identification of the desirable and undesirable plants involved. There are about 100 weeds that commonly occur in the major turfgrasses. These plants can be grouped as desirable grasses, weedy grasses, grass-like weeds, sedges and broadleaf weeds. The following is a brief description of representative plants in each group followed by general suggestions for control.

WEEDY GRASSES	
Goosegrass, <i>Eleusine indica</i>	Annual; clump leaning out from center; stem flattened; center of clump white due to white leaf sheath; leaf sheath at center of plant with a thin green stripe in center; seedhead usually with at least one branch below tip; seeds hang under branch.
Crowfootgrass, <i>Dactyloctenium aegyptium</i>	Annual; clump or running; leaf blade with thin stiff hairs along margin, hairs occur from base to tip or at least over half the length of the blade; ligule a thin transparent membrane seen with magnification; seedhead with all branches at the tip; tip of branch with a small sharp point; seeds hang under branch.
Crabgrass, Five species exist in Turf.	<p>India Crabgrass (<i>Digitaria longiflora</i>) - Annual; running aboveground; usually found in dry sites; no hairs on plant; ligule a thin transparent membrane visible to naked eye, but tiny; leaf blades quite short (<2").</p> <p>Blanket Crabgrass (<i>D. serotina</i>) - Annual; running aboveground, mat-forming; short leaves (<2"), hairy; found in moist to wet sites; same ligule as India crabgrass. Native.</p> <p>Southern Crabgrass (<i>D. ciliaris</i>) - Annual; clumps or loosely running; large leaves (>2"), hairy; found in dry sites; same ligule as India crabgrass; seedhead branches from several points.</p> <p>Tropical Crabgrass (<i>D. bicornis</i>) - Annual; clumps or loosely running; large leaves (>2"), hairy; found in dry sites; same ligule as India crabgrass; seedhead branches all from the same point.</p> <p>Smooth Crabgrass (<i>D. ischaemum</i>) - Annual; clumps or loosely running; large leaves (>2"); few hairs on plant; same ligule as India Crabgrass. Found mainly in north Florida and the panhandle.</p>
Thin or Bull Paspalum, <i>Paspalum setaceum</i>	Perennial; clump leaning out from center; hairy or smooth; hard short knotty root structure; seedheads with usually one or occasionally 2 or 3 branches; tiny seeds hang from lower side of branches.
Tropical Signalgrass (a.k.a. Smallflowered Alexandergrass <i>Urochloa subquadriflora</i>)	Perennial from stolons; leaf blade and sheath hairy; seedheads with two to seven branches or "fingers"; angle of branches resembling a "signal flag"; reproduces by seed and stolons.

LEAVES RESEMBLING GRASS LEAVES WITH PARALLEL VEINATION, BUT LACK A LIGULE AND USUALLY A SHEATH	
Doveweed <i>Murdannia nudiflora</i>	Annual; succulent; loosely running; leaf sheath closed; few hairs at base of blade on margin; tiny purple or blue-purple flower; seeds in tiny round capsules; in dayflower family.
Spreading Dayflower <i>Commelina diffusa</i>	Annual; succulent; leaves broadly lance-shaped, with closed sheaths; sheaths short with a few soft hairs on upper margin; flowers with three blue petals, in leaf-like structure open on the margins.
Annul blueeyed-grass <i>Sisyrinchium rosulatum</i>	A winter annual with flattened leaves that cluster at the base of the plant resembling goosegrass when it is small but germinates in the fall and produces a small blue or purple flower in the spring.

BROADLEAVES	
Match-head, <i>Phyla nodiflora</i>	Perennial; running aboveground; leaves opposite; leaves with teeth on margin; plant gray-green; flowers in heads on long stalks resembling a match; flowers at tip purplish to white.
Erect and Prostrate Spurge, <i>Chamaesyce</i> species	Annual; erect or prostrate branched stem; leaves opposite; base of leaf not equal; sap milky. Large number of species occur in FL.
Chamberbitter, <i>Phyllanthus urinaria</i>	Annual; erect branched stem; leaves alternate; fruit or small round capsules on short-stalks hang under the branches; capsules have warts.
Sagotia Beggarweed, <i>Desmodium triflorum</i>	Perennial; runners aboveground; leaves trifoliolate (three leaflets), alternate; leaflets on short stalks, broader toward tip; flowers small, purple; fruit with 3 to 5 segments.
Rustweed, <i>Polypremum procumbens</i>	Annual or perennial; forming clumps; leaves opposite, quite narrow, needlelike; flowers white, small 4-lobed; fruit dry, indented at tip; leaves turn a rust color, hence the name.
Common Beggarticks, <i>Bidens alba</i>	Annual; erect branched stem, 2 inches to 9 feet tall; leaves opposite, simple (one leaf) on seedlings, young plants and the lower parts of older plants, leaves compound (3 to 9 leaflets) on mature plants; flowers white; fruits long, narrow and stick to clothing; in Sunflower Family.
Brazil Pusley, <i>Richardia brasiliensis</i>	Perennial; branched spreading stem; plant quite hairy; leaves opposite; leaf margin smooth; root thickened, fleshy, with a thin upper portion so that it is easily broken when pulled; flowers white in a dense cluster at stem tips; fruits with stiff hairs.
Florida Pusley, <i>Richardia scabra</i>	Annual; branched spreading stem; plant quite hairy; leaves opposite; leaf margin smooth; root a tap root, thick near plant and tapering downward; flowers white in a dense cluster at stem tips; fruits with bumps.
Florida Betony <i>Stachys floridana</i>	Perennial; branched erect stem from thin white underground runners and fleshy white tubers; leaves opposite, simple, shovel-shaped, toothed, stalked; flowers pinkish-purple; fruit composed of four nutlets.
Pennywort (dollarweed) <i>Hydrocotyle</i> spp.	Perennial from rhizomes, occasionally with tubers; erect long-stalked leaves with scalloped margins; petiole in center of leaf, "umbrella-like," rather than at edge as in Dichondra; found in moist to wet sites; reproduces by seed, rhizomes, and tubers.
Lawn Burweed (Spurweed) <i>Soliva pterosperma</i>	Low-growing, freely branched winter annual. Leaves opposite, sparsely hairy and twice divided into narrow segments or lobes. Flowers small and inconspicuous. Fruits clustered in leaf axils having sharp spines that can cause injury to humans. Reproduces by seed.

BROADLEAVES	
Virginia Buttonweed <i>Diodia virginiana</i>	Spreading perennial herb with hairy branched stems. Leaves opposite, elliptic to lance-shaped, sessile, joined across stem by membrane. Membrane with a few "hair-like" projections. White tubular flowers with four lobes at each leaf axil along the stem. Flower usually with only two sepals. Fruit green, elliptically shaped, hairy, ridged and at each leaf axil. Reproduces by seed, roots, and stem fragments. Favors moist to wet sites.
Old World Diamond-flower <i>Hedyotis corymbosa</i>	Smooth, spreading summer annual with branched stems. Leaves opposite and narrow. Flowers white, usually two or more on long stalks from the tip of a common long stalk. Flowers midsummer until frost. Found in moist areas especially areas that have been disturbed.

PRE-PLANT NONSELECTIVE WEED CONTROL <i>(Refer to Herbicide Label for Specific Use Listing)</i>			
COMMON NAME	TRADE NAME	WEEDS CONTROLLED	COMMENTS
Methyl bromide	Dowfume MC-2 Bromogas Profume Terrogas	Non-selective	Methyl bromide is formulated as liquid gas under pressure that forms a vapor when released. One to 1½ lb material is required per 100 sq.ft. treated soils. Use the higher rate when soils are heavy in texture, wet, or soil temperatures are below 60 F. Soil should be moist but not saturated. Plow soil 8 to 10 inches in depth and release the chemical under a gas proof (plastic) cover. Most other soil pests are also controlled. Grass can be planted 2 to 3 days after application. Methyl bromide is a toxic material used by professional applicators only. Restricted Use Pesticide.
Metham-sodium (metham) Dazomet	Vapam Basamid Granular	Non-selective	A cover is not required but increased control usually results with one. When a cover is not used, cultivate the soil to the desired depth of metham penetration. Soil temperatures should be above 50F before use. Moisten the soil and use 1 to 2 pints of metham product per 100 sq.ft. in 2 to 5 gallons of water. Dazomet rate is 8 to 10 oz product per 100 sq.ft. of prepared soil surface and should immediately be incorporated with a rotary tiller 4 to 8 inches deep and sealed with water at 15 gals. per 100 sq.ft. Immediately irrigate metham to the depth control is desired. If a cover is available, treat the soil in front of a rotary tiller. Cover the soil for 2 days, Planting may take place 2 to 3 weeks after treatment. Aeration may be required by rototilling before planting. Read and follow all label directions. Metham is now a restricted-use-pesticide while Dazomet is not.
1,3-dichloropropene	Telone II	Non-selective	1,3-dichloropropene applied at rates greater than 35 gallons per acre AND covered with a plastic tarp will provide effective control of most annual and perennial weed species.
Glyphosate	Many	Non-selective Non-selective	Glyphosate is applied only to unwanted vegetation and will not control non-germinated seeds, diseases, nematodes, or other pests. Glyphosate (4 lb/gal) is applied at 2 oz. per gallon of water. Wait 2 to 3 weeks after application for regrowth and re-apply. A minimum of 3 applications will be required to control bermudagrass or torpedograss. Do not apply to desirable plants. Same rates as for RoundUp Pro (comparing acid equivalent) - see label for details.
Diquat	Reward Landscape and Aquatic Herbicide	Non-selective	Burn-down of undesirable above-ground grass and broadleaf weed growth. Spot spray using 1-2 qts per 100 gallons of water. For broadcast application, use 1-2 pints per acre. To obtain expected results, a surfactant must be added.
Glufosinate ammonium	Finale	Non-selective	Glufosinate is a nonselective water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds. Glufosinate (1 lb/gal) is applied at 1.5-4.0 oz per gallon of water. Do not apply to desirable plants.

PREEMERGENCE CONTROLS¹ (Refer to Herbicide Label for Specific Species and Use Listing)

Comments. Preemergence herbicides provide 90 to 100 days residual control and require repeat applications for season-long effectiveness. Approximate timings of application for preemergence crabgrass control are: February 1 in south Florida; February 15 in central Florida; and March 1 in north Florida. Goosegrass germinates approximately 3 to 4 weeks later than crabgrass. Adequate soil moisture, both prior to and following application, is necessary to ensure success. Dinitroaniline herbicides (e.g., benefin, oryzalin, pendimethalin, and prodiamine) are not recommended on high traffic areas such as athletic fields, cart paths, par-three tees, and areas not well established. For these high traffic areas with goosegrass, consider using a product containing oxadiazon for annual grass control and simazine for broadleaf weed control. Many herbicides are formulated as "stand alone" products as well as on granules in combination with a dry fertilizer as "weed-and-feed" products.

Grass	Common Name (lbs ai/acre) ²	Trade Name (rate of product/acre)	Weeds Controlled	Comments
Bermudagrass	fenarimol (see comment)	Rubigan 1AS (see comment)	<i>Poa annua</i>	A systemic fungicide that reduces the infestation of <i>Poa annua</i> . Use 3 applications. Treatments should be spaced 10-14 days apart with the third 2 weeks prior to ryegrass overseeding and 30 day interval for <i>Poa trivialis</i> or bentgrass. Use 4 oz/1000 sq. ft. each for 3 applications; or 6 oz/1000 sq. ft. each if 2 applications are used instead of 3. A follow-up application of 2 oz/1000 sq. ft. may be necessary in early January for season-long control where weed pressure is traditionally heavy. Provides little postemergence control. See supplemental label for more information.
	pronamide (1.0 lb)	Kerb 50 W (2.0 lbs)	<i>Poa annua</i> Crabgrass Cool-season grasses	Make application at least 60 days prior to overseeding. Do not apply on or up slope to cool-season turf. Activated charcoal can be used at 2 to 5 lbs/1000 sq. ft. to "deactivate" pronamide when applied closer than 60 days prior to overseeding. Restricted Use Product.
	ethofumesate (1.0 lb)	Prograss 1.5 EC (2.67 qt)	<i>Poa annua</i>	Provides <i>Poa annua</i> control in dormant (full, complete dormancy for > 2 months) bermudagrass overseed with perennial ryegrass. The first application should be 30 to 45 days following overseeding. The second should be 21 to 28 days later. Do not apply after February 1. Not labeled for greens or zoysiagrass.
Bermudagrass St. Augustinegrass Zoysiagrass	oxadiazon (3.0 lbs)	Ronstar 2G (150 lbs) Ronstar 50WP	Same as for benefin, goosegrass	For use on Bermuda, St. Augustine, & zoysiagrasses only. Do not apply to wet turf or to golf greens. NOT FOR USE ON HOME LAWNS. Ronstar 50WP can be used only on dormant turf or excessive phytotoxicity will result. Thoroughly irrigate following application to increase effectiveness. Safest preemergence herbicide on newly sprigged or high traffic areas. A combination of oxadiazon plus benefin on a 38% ureaformaldehyde nitrogen fertilizer is available as Regal Star.

Grass	Common Name (lbs ai/acre) ²	Trade Name (rate of product/acre)	Weeds Controlled	Comments
Bahigrass Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	benefin (3.0 lbs)	Balan 2.5G (120 lbs) 2.5 Benefin G (120 lbs)	Crabgrass, crowfootgrass, <i>Poa annua</i> , sandbur, some selected broadleaves.	Apply only to well-established turf before annual grass weed seed germination. For continued weed control, a second application 3 months after the initial is required. For <i>Poa annua</i> control, use full rate in September. Minimum 3 month waiting period is required before reseeding. Read the label for irrigations requirements to activate the herbicide. DO NOT APPLY TO IMMATURE TURF, desirable overseeding, or on golf greens.
	benefin + trifluralin (2.0 lbs + 1.0 lb)	Team 2G (150 lbs)	Same as for benefin	Same as for benefin. For use by professional applicators only. Good for use in mixed stands containing cool and warm-season turfgrasses.
	bensulide (7.5 - 12.5lbs)	Betasan 3.6G (209-348 lbs) Bensumec, 4LF (1 ⁷ / ₈ -3 ¹ / ₈ gal) Pre-San 7G (107-179 lbs) Pre-San 12.5G (60-100 lbs) ProTurf Weedgrass Preventer 8.5G (88-147 lbs)	Same as for benefin	Same as for benefin. Safe on overseeded areas and golf greens. If use on putting greens, apply at least 90 days before overseeding. Bensumec 4LF can be applied to dichondra lawns at the time of seeding or any time thereafter.
	DCPA (10.5 lbs)	Dacthal W-75 (14 lbs)	Same as for benefin, prostrate spurge	Same as for benefin. May be applied to seedlings when they reach 1 to 2 inches in height. A repeat application at a half rate is needed 60 days after the first to extend the control period.
	dithiopyr (0.38 - 0.50 lbs)	Dimension 1EC (0.5 gal) Dimension Ultra 2SC (24 oz) Dimension 40 WP (0.95 lbs)	Same as for benefin, goosegrass	Same as for benefin. Do not use within 45 days of seeding or sprigging. A total of 1 ¹ / ₂ lb ai/A is allowed yearly but not to exceed ¹ / ₂ lb ai/A per application. Preemergence crabgrass control may require a second application 60-90 days after initial application. Postemergence activity on 2-3 leaf stage crabgrass.
	oryzalin (1.5 lbs)	Surflan 4AS (1.5 qts)	Same as for benefin, goosegrass	Same as for benefin. Use a 1 ¹ / ₂ + 1 ¹ / ₂ lb ai/A split application approximately 90 days apart for best results. Longest period (21 days) for preemergence herbicide before required activation by rainfall or irrigation. Spring application on overseeded, cool-season grasses may prematurely thin them.
	oryzalin + benefin (1.5 lbs + 1.5 lbs)	XL 2G (150 lbs)	Same as for benefin, goosegrass	Same as for benefin

Grass	Common Name (lbs ai/acre) ²	Trade Name (rate of product/acre)	Weeds Controlled	Comments
Bahigrass Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	pendimethalin (2.75 lbs)	PRE-M 60DG Pendulum 60 DG (2.5 - 5.0 lbs) PRE-M 3.3 EC Pendulum 3.3 EC (4.2 - 7.2 pts) PRE-M/Fertilizer (check label) Pendulum AquaCap 3.8 ME (4.2 - 6.3 pts)	Same as for benefin, goosegrass, oxalis, speedwell	Same as for benefin. For use by professional applicators only. A split application of 1½ to 2.0 lb ai/A before weed seed germination followed by a 1 to 1½ lb ai/A application 90 days later provides better season-long control, especially when heavy weed pressure is expected. Check the product label for registration on golf greens. Spring application on overseeded, cool-season grasses may prematurely thin them. Sequential applications of Pendulum can be made at 60 to 90 days after initial application for longer control.
	S-metolachlor (1.1 - 2.2 lbs)	Pennant Magnum 7.62L (1.3 - 2.6 pts)	Yellow nutsedge, annual sedge, sprangletop, some annual grass suppression	For use on golf course fairways, sod farms, and commercial lawns. The higher rate will be necessary for turf grown on high organic (i.e., muck) soils. For commercial St. Augustinegrass sod production, do not use more than once every 6 weeks and do not apply more than 4.2 pts./A/yr. Tank mixing with atrazine will increase the weed control spectrum. Do not use Pennant Magnum on golf greens, tees, or aprons or within 4 months of overseeding. Irrigate within 7 days after application.
	prodiamine (0.5 - 1.5 lbs)	Barricade 65WG (0.5 - 2.3 lbs) Barricade 4FL (0.6 - 3 pts) RegalKade (check label) ProClipse (check label)	Crabgrass, crowfootgrass, <i>Poa annua</i> , goosegrass, Signalgrass - broadleaf, spurge, Pusley - Florida, etc.	Apply to well-established turf before annual weed seed germination. Barricade may be applied as a single application or in sequential applications to control weeds germinating throughout the year. This is rate dependent. Split applications of low rates may be made at 60 to 90 day intervals. Do not apply to overseeded turf within 60 days after seeding. May be used on newly sprigged bermudagrass at rates not to exceed 0.8 lbs / A. RegalKade formulations are on dry fertilizer carriers.
	isoxaben (1.0 lb)	Gallery 75W (1.33 lbs)	Broadleaves	Controls broadleaf weeds. Tank mix with another preemergence herbicide for grass weed control. In order to activate the material, ½" water is needed following application. Not labeled for golf greens. Do not reseed until 30 days after application. Do not apply to newly seeded turf until it has been mowed 3 times.

Grass	Common Name (lbs ai/acre) ²	Trade Name (rate of product/acre)	Weeds Controlled	Comments
Bermudagrass Centipedegrass St. Augustinegrass Bahia grass	napropamide (2.0 lbs)	Devrinol 50WP (4.0 lbs) Devrinol 2G (100 lbs) Devrinol 5G (40 lbs) Ornamental Herb. 5G (80-120 lbs)	Same as for benefin	Do not apply to immature turf. A second application 8 to 10 weeks after the first is suggested. Not recommended for putting greens. Use the reduced rates for turf maintained at lower mowing heights. Irrigate after application. Do not reseed or overseed within six months after application.
Centipedegrass St. Augustinegrass Zoysiagrass	atrazine/simazine (2.0 lbs-sandy soil) (4.0 lbs-muck soil)	Atrazine Aatrex 4L, 90DG, 80W; Purge Simazine Princep Liquid + others	Same as for benefin plus pennywort (dollarweed), henbit, chickweed, lawn burweed (or spurweed) and some annual sedges. Perennial broadleaf weeds such as will garlic, dock and others usually escape.	SEE LABEL RESTRICTIONS! Apply to centipedegrass, St. Augustinegrass, and zoysiagrass only. Will provide good to excellent weed control with a minimum of growth retardation to newly sprigged, sodded, or plugged turf areas. Effectiveness will be reduced as weeds germinate and mature. Two applications are allowed per year. Pennywort is easiest to control with a late fall and/or early winter application followed by a repeat application 4 to 6 weeks later. Do not apply within the root zone of ornamentals. Do not exceed 1 lb ai/A on newly sprigged turfgrass. Atrazine is a Restricted Use Pesticide.
Centipedegrass Perennial Ryegrass St. Augustinegrass (Sod)	mesotrione (0.125 - 0.25 lb)	Tenacity 4L (4 - 8 fl. oz)	Pre- and postemergence control of annual grass and broadleaf weeds. See label.	Tenacity may be used for weed control in centipedegrass, dormant bermudagrass, and perennial ryegrass in commercial sites. Use sites include non-crop areas: golf courses, sod farms, athletic fields, parks, commercial properties, cemeteries, and airports. Do not use on golf course putting greens and maintain a five foot buffer between treated areas and putting greens. Tenacity may be used on St. Augustinegrass for sod production only. If applied preemergence combine with a preemergence herbicide for extended control of key grass species such as crabgrass and goosegrass. For postemergence applications add a non-ionic surfactant. Postemergence control may require a repeat application 2 to 3 weeks after the initial treatment.

Grass	Common Name (lbs ai/acre) ²	Trade Name (rate of product/acre)	Weeds Controlled	Comments
Bahagrass Bermudagrass Centipedegrass Seashore paspalum St. Augustinegrass Zoysiagrass	proflam + sulfentrazone (See comments)	Echelon (See comments)	Crabgrass, goosegrass, <i>Poa annua</i> , crowfootgrass, signalgrass, Florida pusley, yellow nutsedge, annual Kyllinga, annual sedge	Bahagrass, centipedegrass, seashore paspalum, and zoysiagrass rate: 0.57 - 0.75 lb ai/A, 18-24 fl oz/A per application not to exceed 1.125 lb ai/A, 36 fl oz/A per calendar year. St. Augustinegrass rate: 0.57 lb ai/A (18 fl oz/A) followed by 0.57 lb ai/A (18 fl oz/A) 45 to 60 days later. Temporary discoloration may occur. Bermudagrass rate: 0.75 - 1.125 lb ai/A, 24 - 36 fl oz/A.
	dimethenamid-P (1.0 - 1.5)	Tower 21 - 32 oz	Crabgrass, goosegrass Broadleaves	For use on golf courses and landscaped ornamental areas. DO NOT apply to residential turfgrass, sod farms, or any other improved or unimproved maintained turfgrass areas.
	indaziflam (0.03 - 0.06)	Specticle 2.5 - 5.0 oz	Annual grasses Annual sedges Broadleaves	Do NOT apply to newly seeded turf (> one year old). Do NOT apply to turf that is showing signs of stress. Do NOT apply to golf course putting greens. Rate varies with turfgrass species, region of the state, and time of year. This product provides very long (up to 8 months) residual control of crabgrass and goosegrass.
Bermudagrass Centipedegrass St. Augustinegrass Seashore paspalum Zoysiagrass	dimethenamid-P + pendimethalin (1.75 - 3.5)	Freehand 100 - 200 lbs	Annual bluegrass Crabgrass Doveweed Goosegrass Spurge	Labeled for use on golf courses, sod farms, commercial sites, athletic and sports fields, residential turf, and recreational and parks.
Seashore Paspalum	proflam (0.5 - 1.5 lbs)	Barricade 65WG (0.5 - 2.3 lbs) Barricade 4FL (0.6 - 3 pts) RegalKade (check label) ProClipse (check label)	Crabgrass, crowfootgrass, <i>Poa annua</i> , goosegrass, Signalgrass - broadleaf, spurge, Pusley - Florida, etc.	Apply to well-established turf before annual weed seed germination. Barricade may be applied as a single application or in sequential applications to control weeds germinating throughout the year. This is rate dependent. Split applications of low rates may be made at 60 to 90 day intervals. Do not apply to overseeded turf within 60 days after seeding. May be used on newly sprigged bermudagrass at rates not to exceed 0.8 lbs / A. RegalKade formulations are on dry fertilizer carriers.
	dithiopyr (0.38 - 0.50 lb)	Dimension 1EC (0.5 gal) Dimension Ultra 2SC (24 oz) Dimension 40 WP (0.95 lbs)	Same as for benefin, goosegrass	Same as for benefin. Do not use within 45 days of seeding or sprigging. A total of 1½ lb ai/A is allowed yearly but not to exceed ½ lb ai/A per application. Preemergence crabgrass control may require a second application 60-90 days after initial application. Postemergence activity on 2-3 leaf stage crabgrass.

Grass	Common Name (lbs ai/acre) ²	Trade Name (rate of product/acre)	Weeds Controlled	Comments
Seashore paspalum	oxadiazon (3.0 lbs)	Ronstar 2G (150 lbs)	Same as for benefin, goosegrass	Make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectional injury.
	pendimethalin	Check specific labels.	Same as for benefin, goosegrass, oxalis, speedwell	A split application of 1½ to 2.0 lb ai/A before weed seed germination followed by a 1 to 1½ lb ai/A application 90 days later provides better season-long control, especially when heavy weed pressure is expected.

¹Presence of a herbicide in this listing does not constitute a recommendation. Trade names are used with the understanding that no endorsement is intended or no criticism is implied of similar products which are not mentioned. All chemicals should be used in accordance with the manufacturer's instructions.

²All herbicide rates are active ingredient rates per acre. For product rates for formulations not listed, check the label included with every herbicide container.

Preemergence herbicides for putting greens (*Refer to Herbicide Label for Specific Turf Species and Use Listing*).

Trade Names	Ingredients	Manufacturer/Distributor
Bensumec 4LF	bensulide	PBI/Gordon Corp.
Pre-San	bensulide	PBI/Gordon Corp.
Weedgrass Preventer	bensulide	Andersons
Goosegrass/Crabgrass Control	bensulide + oxadiazon	Andersons
Southern Weedgrass Control	pendimethalin	Andersons
Betasan	bensulide	Gowan
Kerb	pronamide	Dow AgroSciences
Rubigan	fenarimol	Gowan

Turfgrass Tolerance to Preemergence Herbicides (Refer to Herbicide Label for Specific Species Listing)¹							
Herbicide	Bahiagrass	Bermudagrass	Centipedegrass	Seashore Paspalum	St. Augustine	Zoysiagrass	Overseed Rye/Blends
atrazine (Aatrex)	NR ²	NR	S	NR	S	I-S	D
benefin (Balan)	S	S	S	NR	S	S	NR
benefin+oryzalin (XL)	S	S	S	NR	S	S	NR
benefin+trifluralin (Team)	S	S	S	NR	S	S	NR
bensulide (Betasan, Bensumec, PreSan)	S	S	S	NR	S	S	I-S
bensulide+oxadiazon	NR	S	NR	NR	NR	S	NR
DCPA (Dacthal)	S	S	S	NR	S	S	NR
dithiopyr (Dimension)	S	S	S	S	S	S	I
dimethenamid-P (Tower)	S	S	S	S	S	S	S
dimethenamid+pendimethalin (Freehand)	S	S	S	S	S	S	S
ethofumesate ³ (Prograss)	NR	S-dormant	NR	NR	I	NR	S(D)
fenarimol (Rubigan)	NR	S	NR	NR	NR	NR	S
indaziflam (Specticle)	S	S	S	S	S	S	D
isoxaben (Gallery)	S	S	S	NR	S	S	NR
mesotrione (Tenacity)	NR	S-dormant	S	NR	I	NR	S(D)
metolachlor (Pennant Magnum)	S	S	S	NR	S	S	D
napropamide (Devrinol)	S	S	S	NR	S	NR	NR
oryzalin (Surflan)	S	S	S	S	S	S	NR
oxadiazon (Ronstar)	NR	S	NR	S	S	S	I
pendimethalin (Pendulum)	S	S	S	S	S	S	I
prodiamine (Barricade)	S	S	S	S	S	S	I
prodiamine + sulfentrazone (Echelon)	S	S	S	S	NR	S	S
pronamide (Kerb)	NR	S	NR	NR	NR	NR	D
simazine (Princep Liquid)	NR	I	S	NR	S	S	D

¹Non golf green only.²S=Safe at labeled rates on mature, healthy turf; I=Intermediate safety - may cause slight damage to mature, healthy turf. Use only one-half the normal rate when temperatures are hot (>85 F) or if the turf is under water stress; D=Damaging - do not use; NR=Not Registered for use on this turf species.³Ethofumesate is labeled only for Dormant bermudagrass overseeded with perennial ryegrass.

Preemergence Herbicide Efficacy Ratings																							
	atrazine	benefin	benefin + oryzalin	benefin + trifluralin	bensulide	bensulide + oxadiazon	DCPA	dithiopyr	dimethenamid-P	ethofumesate	fenarimol	indaziflam	isoxaben	mesotrione	metolachlor	napropamide	oryzalin	oxadiazon	pendimethalin	proflaminate	proflaminate + sulfentrazone	pronamide	simazine
PERENNIAL WEEDS																							
bahiagrass	F	P	P	P	P	P	P	P	P	P	P	P	P		P	P	P	P	P	P	P	P	P
bermudagrass	P	P	P	P	P	P	P	P	P	P	P	P	P		P	P	P	P	P	P	P	P	P
dallisgrass	P	P	P	P	P	P	P	P	P	P	P		P		P	P	P	P	P	P	P	P	P
nutsedge, purple	P	P	P	P	P	P	P	P	P	P	P	P	P		P-F	P	P	P	P	P		P	P
nutsedge, yellow	P	P	P	P	P	P	P	P	F	P	P	P	P		G	P	P	P	P	P		P	P
tall fescue	F	P	P	P	P	P	P	P	P	P	P		P		P	P	P	P	P	P		G	F
wild garlic/onion	P	P	P	P	P	P	P	P	P	P	P	P	P		P	P	P	P	P	P		P	P
ANNUAL GRASSES																							
<i>Poa annua</i>	E	E	E	E	F	F-G	G	G-E	G	G-E	G	E	P-F	F	G	G	G	G	G	E	E	E	E
crabgrass	F	E	E	E	E	E	G	G-E	G	G	P	E	P	E	G	G-E	E	E	E	E	E	F	F
crowfootgrass	P-F	G	G	G	G	G		G			P		P		F	F	G	G	G	G	G		
goosegrass	P	F	G	G	F	G	F	F-G	G		P	E	P		F	G	F-G	E	F-G	F-G	F-G	F	F
sandbur	P	F	F-G	G	G	F-G	F	F	G		P		P		F	F	G	F	G	G	G		
BROADLEAF WEEDS																							
chamberbitter (niruri)	G										P		G										
common chickweed	E	G	G	G	P	P	E	G		G	P	G	E	E	F	E	G	P	G	G		E	E
corn speedwell	E	E	E	E	P	P	G	G			P		G-E		G	E	P	G	E	E		G	G
cudweed	E	P	P	P		P					P	G	G			G	P	P	G	P		P	
dandelion	F	P	P	P	P	P					P	G	G				P	P	P	P		P	P
dichondra	G	P	P	P	P	P					P						P	P	P	P		P	P
docks	G	P	P	P	P	P					P						P	P	P	P		P	P
doveweed	G	P	P	P	P	P			F	P	P	F					P	P	P	P		P	G
Florida betony	E	P	P	P	P	P					P						P	P	P	P		P	P

Preemergence Herbicide Efficacy Ratings																							
	atrazine	benefin	benefin + oryzalin	benefin + trifluralin	bensulide	bensulide + oxadiazon	DCPA	dithiopyr	dimethenamid-P	ethofumesate	fenarimol	indaziflam	isoxaben	mesotrione	metolachlor	napropamide	oryzalin	oxadiazon	pendimethalin	proflaminate	proflaminate + sulfentrazone	pronamide	simazine
ground ivy		P	P	P	P	P					P						P	P	P	P		P	P
BROADLEAF WEEDS																							
henbit	E	G	G	G	P	P	F	G			P	G	G			P	G	P	G	G		F	E
hop clovers	E	P	G	G	P	F					P			G			F	G	G	P		P	E
knotweed	E		G	G	G	G		G			P		G			G	F	G	G			G	G
lespedeza	E	P	P	P				G			P						P		P	P			E
mallow		P	P	P	P	P					P		G				P	P	P	P		F	P
mock strawberry		P	P	P	P	P					P						P	P	P	P		P	P
mouseear chickweed	E	E	E	E	P	P	G	G			P		G	E			P	P	G	G		G	G
mugwort		P	P	P	P	P					P						P	P	P	P		P	P
mustards	E				G	G-E	P	G			P		G					E				F	F
parsley piert	E	P	P	P	E	G-E		G			P							G	P	P		P	G
pennywort (dollarweed)	E	P	P	P	P	P					P		G				P	P	P	P		P	P
plantains	G	P	P	P	P	P					P	G	G				P	P	P	P		P	P
spurges	E	P	F	F	P	P	F	G	G		P		G		F	P	F	P	F	F		P	G
spurweed (burweed)	E	P	P	P	P	P	P				P		E			E	P	P	P	G		P	E
VA buttonweed		P	P	P	P	P					P						P	P	P	P		P	P
violets		P	P	P	P	P					P						P	P	P	P		P	P
white clover	E	P	P	P	P	P					P		G				P	P	P	P		P	G
yellow woodsorrel (<i>Oxalis</i>)	E	P	F-G	F-G	P	F		G			P	G	G		P	G	F	G	F-G	F		P	P

Key to response symbols: E = Excellent control (90 to 100%), G = Good control (80 to 89%), F = Fair control (70 to 79%), P = Poor control (< 70%). A blank space indicates weed response is not known.

POSTEMERGENCE CONTROLS (Refer to Herbicide Label for Specific Turf Species and Use Listing)

Comments: Active only on emerged, visible weeds. Best results occur when weeds are young. Temperatures above 85-90° may result in phytotoxicity (yellowing) to the turf. Repeat applications may be required for acceptable control. These should be timed 10 to 14 days apart. Do not mow within 48 hrs after application for most chemicals. Most postemergence herbicides require the use of a spreader-sticker, adjuvant, crop oil, or wetting agent. Read the label before adding these as many herbicides are pre-packaged with them already added.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass	diclofop-methyl (0.75 - 1.0 lb)	Illoxan 3EC (1.0 - 1.4 qts)	Goosegrass	Do not apply more than one (1) fluid ounce per 1,000 sq. ft. per treatment and not more than a total of 1.5 fluid ounces per 1,000 sq. ft. per year. Young goosegrass plants are easiest to control. The high rate is needed for older plants. Larger, mature goosegrass will not be adequately controlled. Do not mow 24-36 hours after applying. Control takes 2-3 weeks. Treat only well established and actively growing turf. Wait 6 weeks before overseeding after the last application. DO NOT tank mix Illoxan with any other pesticide or liquid fertilizer as reduced goosegrass control may occur. Restricted Use Pesticide.
	metribuzin (0.25 lb)	Sencor 75DF (0.33 lb)	Goosegrass, some broadleaf weeds	
	pronamide (1.50 lbs)	Kerb 50W (3.0 lbs)	<i>Poa annua</i> , ryegrass clumps	Do not apply on or up-slope to desirable overseeded turf as pronamide may run. Time required for control increases as weeds mature, therefore apply in late fall for optimum results. If applied in spring, 3 to 4 weeks are required for control. Restricted Use Pesticide.
	simazine (1.0 lb)	Princep Liquid (1 qt)	<i>Poa annua</i> , most annual winter broadleaf weeds	Do not exceed use rates. For winter annual weed control, apply 1 qt/A in early fall (after Oct. 15) and repeat in early winter. Some yellowing of bermudagrass may result. Do not apply on or up slope to desirable overseeded turf &/or golf greens. Do not use on bermudagrass during spring 'green-up' or summer. Resistance to this herbicide has been noted where continued use without rotation occurs.
	rimsulfuron (0.008 - 0.031 lb)	TranXit (0.5 - 2.0 oz)	Ryegrass Removal <i>Poa annua</i>	REMOVAL OF OVERSEED: Apply recommended rate in the spring months 3 to 4 weeks before desired date for overseed removal. <i>POA ANNUA</i> CONTROL: Apply recommended rate within the period of 10 to 14 days prior to overseeding perennial ryegrass and/or <i>Poa trivialis</i> .

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass overseeded with Perennial ryegrass	bispyribac-sodium (0.02 - 0.06 lb)	Velocity (2.0 - 6.0 oz)	<i>Poa annua</i>	Apply two or three times on a 14 to 21 day interval at 1.33 oz/A for control of heavy <i>Poa annua</i> infestations. Use season is January 1 to April 15. Apply to actively growing <i>Poa annua</i> when it first begins to flower. Do not apply when temperatures are below 55 °F.
Bahigrass Bermudagrass Zoysiagrass Ryegrass	2,4-D Amine (1.0 lb) See product label.	Several Brands	Many broadleaf weeds including matchweed, pennywort, (dollarweed), wild garlic/onion, clover, plantains.	Apply when weeds are young and actively growing. Repeat application in 10 to 14 days may be necessary for complete control. Use lower rates (0.5 lb ai/A) on `Tifgreen' and `Tifdwarf' Bermudagrass. Amine formulations should be used as high volatile ester formulations have drift and volatility problems. For hard-to-control perennial broadleaf weeds, formulations containing dicamba will increase control. One-half rate should be used on centipede grass and carpet grass.
	2,4-D + 2,4-DP See product label.			
	dicamba (0.125 - 0.25 lb) See product label.	Banvel 4S (0.25 - 0.50 pts) plus others	White clover, spurge, woodsorrel, dichondra	Avoid drift. Do not apply within the root zone of ornamentals. Repeat applications 10 to 14 days apart may be needed for complete control but may also result in some turf injury. Check label for use on greens.
	dicamba (0.125 lb) + 2,4-D, MCPP, MCPA, and/or 2,4-DP (0.50 - 0.75 lb)	Several brands contain these mixtures. See product label for specific rates.	Same as for dicamba, also matchweed, pennywort.	Same as for dicamba. Refer to product label for rates as herbicide ratios vary depending on brands. Use only on actively growing, non-stressed turf. Check label for use on golf greens
	triclopyr + clopyralid (0.28 - 0.56 lb) + (0.09 - 0.18 lb)	Confront (1-2 pts)	Black medic White clover Lawn burweed Woodsorrel Virginia buttonweed	Avoid drift. Do not apply to exposed roots of trees or shrubs. Do not use on golf greens.
Bahigrass Bermudagrass Centipede grass Seashore paspalum Zoysiagrass	proflam + sulfentrazone	Echelon (See comments)	Annual grasses and some broadleaves	Bahigrass, centipede grass, seashore paspalum, and zoysiagrass rate: 0.57 - 0.75 lb ai/A, 18-24 fl oz/A per application not to exceed 1.125 lb ai/A, 36 fl oz/A per calendar year. Bermudagrass Rate: 0.75 - 1.125 lb ai/A, 24 - 36 fl oz/A.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Centipedegrass St. Augustinegrass Zoysiagrass	atrazine/simazine (1.0 - 2.0 lbs)	Several Brands. Read the label for rates	Many broadleaf weeds including matchweed, oxalis, pennywort, Florida betony and some annual sedges.	SEE LABEL RESTRICTIONS: For hard to control weeds, make the first application in late winter and follow with another 4 to 6 weeks later. If weeds persist, follow atrazine applications with dicamba in 4 to 6 weeks. Some turf injury can be expected with this. Two applications of atrazine are allowed per year. Effectiveness will be reduced as weeds mature. Do not apply within the root zone of ornamentals.
Centipedegrass St. Augustinegrass Zoysiagrass	dicamba (0.125 - 0.25 lb)	Banvel 4S (0.25 - 0.50 pts)	White clover, spurge, woodsorrel	Avoid drift. Do not apply within the root zone of ornamentals. Use low rates on St. Augustinegrass. Treat when temperatures are ≤80 F to minimize turf damage.
	dicamba + 2,4-D, 2,4-DP, MCPA, and/or MCPP (0.125 + 0.25 + 0.50 lb)	Several brands contain these mixtures	White clover, spurge, woodsorrel, pennywort	Observe same precaution as dicamba above. Refer to product label for rates. A second application on centipedegrass 7-14 days later may be needed. Use low rates on St. Augustinegrass. A tank mix of atrazine at 1 lb ai/A + 2,4-D & dicamba at 0.2 lb ai/A each provides good control with minimum turf damage when temperatures are ≤80 F.
	bromoxynil (0.375 - 0.50 lb)	Buctril 2L (1.0 - 2.0 pts)	Many young broadleaf weeds	Labeled only for non-residential turf, seed and sod production. Contact herbicide. Safe on seedling or sprigged turf. Tank mixing with 2,4-D, dicamba, &/or MCPP will provide increased control but should be used only on established turf. May also be used on Bermudagrass, Bentgrass, and Ryegrass. Restricted Use Pesticide.
Bahiagrass Bermudagrass St. Augustinegrass Zoysiagrass	bromoxynil (0.25 - 0.50 lb)	Buctril 2L (1.0 - 2.0 pts)	Many young broadleaf weeds	Labeled only for non-residential turf, seed and sod production. Safe on seedling or sprigged turf. Tank mixing with 2,4-D, dicamba, &/or MCPP will provide increased control but should be used only on established turf.
Centipedegrass	sethoxydim (0.33 lb)	Segment 1.0L (2.25 pts)	Crabgrass, goosegrass and other annual grasses	Apply before weeds mature. Repeat applications are necessary to suppress Bermudagrass or Bahiagrass. Safe on centipedegrass seedlings after the third mowing.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Centipedegrass	mesotrione (0.156 - 0.25 lb)	Tenacity 4L (5.0 - 8.0 fl oz)	Crabgrass, goosegrass, Oxalis, ground ivy, yellow nutsedge, and other grassy and broadleaf weeds.	For use on SOD FARMS ONLY . Efficacious as an early post- and preemergence (combined) application. Apply to young weeds. Addition of a non-ionic surfactant required. A repeat application within 3-4 weeks may be required for more mature weeds.
	clethodim (0.125 - 0.25 lb)	Envoy (17 - 34 fl. oz.)	Common Bermudagrass control	For use on SOD FARMS ONLY . Do not apply to centipedegrass being grown for seed. Do not apply ENVOY Herbicide until 3 weeks after 100% greenup of centipedegrass in spring. Two Applications will likely be necessary for control. Second application should not be made until 1-2 inches of new growth of bermudagrass is observed (approximately 3-4 weeks after first application). Use higher labeled rates for more established bermudagrass. Do not treat or allow drift of ENVOY Herbicide onto other turf species, or damage may result. Do not apply ENVOY Herbicide if rainfall is expected within one hour after application. Avoid mowing sod for one week before and after application. At rates above 17 oz./A, ENVOY Herbicide can cause temporary centipedegrass injury when tank-mixed with crop oil concentrate.
Bahigrass Bermudagrass	hexazinone (2.0 - 6.0 lbs)	Velpar 2L (1.0 - 3.0 gal)	Smutgrass, some broadleaf weeds	Labeled only for non-cropland turf such as roadsides, railroads, industrial sites, and underneath utility lines. Soil moisture must be present for herbicide activity. Best control and least turf damage follows late winter application. Do not apply after April. Temporary turf discoloration can be expected following treatment, especially to Bahigrass. Do not use underneath desirable shrubs or trees. Read and follow all label directions before use.
	sulfometuron (0.05 -0.19 lb)	Oust 75DG (1.0 - 2.0 oz)	Fescue, Broadleaf weeds	Highway roadside use only. Use in spring (1 to 3 oz/a) 30 days after green-up or 1 to 2 weeks after mowing. Also used in late fall to early winter (1 to 4 oz/a). Often tank-mixed with MSMA (3 to 4 lbs ai/a) for broader spectrum of weed control in bermudagrass.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bahiagrass Bermudagrass	sulfosulfuron (0.5 - 1.5 oz)	Outrider (0.75 - 2.0 oz)	Johnsongrass Purple Nutsedge Yellow Nutsedge Horseweed Vaseygrass Chickweed	Highway roadside use only.
St. Augustinegrass	ethofumesate (3.0 lbs)	Prograss 1.5EC (2 gal)	Common Bermudagrass control or suppression	Timing is critical. Spring applications should start in late Feb. (South FL) or early (central FL) to mid (North FL and Northward) March. Repeat in 30 days. Tank mixing with atrazine or simazine at 2 lb ai/A significantly increases suppression. For repeat applications, reduce atrazine or simazine to 1 lb ai/a. Temporary St. Augustinegrass stunting may result. Do not overlap.
	mesotrione (0.125 lb)	Tenacity 4L (4 fl oz)	Crabgrass, goosegrass, Oxalis, ground ivy, yellow nutsedge, and other grassy and broadleaf weeds.	FOR SOD PRODUCTION ONLY. Efficacious as an early post- and preemergence (combined) application. Apply to young weeds. Addition of a non-ionic surfactant required. A repeat application within 3-4 weeks may be required for more mature weeds. Some temporary discoloration (whitening) may occur.
Zoysiagrass	fenoxaprop (0.12 - 0.35 lb)	Acclaim 1EC (15 - 45 oz)	Annual grass weeds, Bermudagrass suppression	Young, actively growing weeds are easiest to control. Do not apply to moisture- or heat-stressed turf. Repeat application in 2 to 3 weeks may be required for complete control. Do not mow for at least 24 hrs after application.
	fluazifop (0.047 - 0.078 lb)	Fusilade II (2EC) (3 - 4 oz)	Annual grasses Common Bermudagrass Suppression	Add nonionic surfactant. Repeat application in 4 weeks. Minor, short-term turf phytotoxicity may occur, especially when applied during hot, dry weather.
Bermudagrass (Common & Hybrid) Zoysiagrass	carfentrazone + MCPA + MCPP + dicamba	Power Zone (2 - 4 pts)l	Broadleaf weeds (see label)	Carfentrazone combinations are fast-acting often exhibiting evidence of activity within hours.
	carfentrazone + 2,4-D + MCPP + dicamba	Speed Zone (2 - 4 pts)	Broadleaf weeds (see label)	Carfentrazone combinations are fast-acting often exhibiting evidence of activity within hours.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass (Common & Hybrid) Zoysiagrass	foramsulfuron (0.013 to 0.026 lbs)	Revolver Broadcast: (5.5 - 26.2 oz) Spot Treatment: (0.5 - 2.0 fl. oz /gal)	Ryegrass Removal <i>Poa annua</i> Goosegrass	Apply when bermudagrass has resumed active growth and removal of ryegrass is desirable. Speed of ryegrass removal depends on temperature (above 70 F, removal can occur in 1 week).
	quinclorac (0.25 - 0.75 lbs ae/A)	Drive XLR8 (64 fl oz)	Torpedograss Crabgrass	For best torpedograss control, make split applications at 21 day intervals.
	quinclorac + dicamba + MCPP	Onetime Herbicide (64 fl oz)	Torpedograss Crabgrass Broadleaf weeds (see label)	
	trifloxysulfuron (0.00469 - 0.02 lb)	Monument 75WG (0.1 - 0.53 oz)	<i>Poa annua</i> Perennial ryegrass Nutsedge and Kyllinga Broadleaf weeds (see label)	0.1 to 0.35 oz/Ac for perennial ryegrass and <i>Poa trivialis</i> removal. 0.35 to 0.53 oz/Ac to control <i>Poa annua</i> , purple and yellow nutsedge, various broadleaf weeds and grasses in established turf (see label). Use rate of 0.53 oz/Ac for broadleaf signalgrass and suppression of torpedograss, Virginia buttonweed (seedlings), bahiagrass, dallisgrass, and crabgrass. Buffer water to pH 7 or less. Speed of ryegrass/ <i>Poa annua</i> removal depends on temperature. Allow at least 3 weeks between last application and overseeding with cool season grasses for winter cover. Weed death may take 2 to 4 weeks or more under cool conditions. Lightly irrigate 4 hours or more after application to minimize tracking and movement if applications are made next to non-labeled cool-season turfgrass species. Multiple applications needed for hard-to-control weeds.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bahia grass Bermudagrass Centipedegrass Seashore paspalum St. Augustinegrass Zoysiagrass Perennial Ryegrass	clopyralid (0.1 - 0.25 lb)	Lontrel T & O (1/4 - 2/3 pt)	Broadleaf weeds including black medic, clover, and plantain.	Maximum use rate in Florida is 2/3 pt. per acre per growing season. Clopyralid containing products may NOT be used on residential turf.
	fluroxypyr (0.125 - 0.25 lb)	Spotlight (0.66 - 1.3 pts)	Clover and other broadleaf weeds (see label).	DO NOT APPLY TO ST. AUGUSTINEGRASS IN FLORIDA. Approved tank mixtures provide improved weed control spectrum (see label). Use low rates on bermudagrass as a spot treatment.
	carfentrazone + 2,4-D + MCP + dicamba	Speed Zone - Southern (1.5 - 4 pts)	Broadleaf Weeds (see label)	Carfentrazone combinations are fast-acting exhibiting evidence of activity within hours. Do Not Apply to 'Floritam' or 'Bitterblue' varieties. Do Not Apply to St. Augustinegrass under shade, disease and/or moisture stress. During higher temperatures, lower rates are recommended.
	sulfentrazone (0.25 - 0.375 lb)	Dismiss (8.0 - 12.0 fl oz) Spartan (sod)	Yellow Nutsedge Purple Nutsedge Kyllinga sps. Goosegrass Selected broadleaves	May cause temporary discoloration in bahiagrass, St. Augustinegrass, and zoysiagrass. Do not apply to weakened turfgrass.
Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	metsulfuron (0.0156 - 0.0625 lb)	Manor (0.25 - 1.0 oz) Blade (0.25 - 1.0 oz)	Safely removes unwanted bahiagrass from desirable turf species. Selected broadleaves.	Do not expect to see results for two to three weeks after application. Will require repeat application(s).
	pyraflufen-ethyl (0.001 - 0.006 lb)	Octane 2% SC (0.7 - 4.0 fl oz/A)	Broadleaf Weeds (see label for Weeds Controlled)	Use in tank mix combinations with registered herbicides for control of annual and perennial broadleaf weeds or as stand-alone for the control of seedling, non-mature winter and summer annual weeds and/or for temporary burn-down of weeds. PPO herbicide with symptoms often visible in 24 to 48 hours. Not for use on golf course greens or tees.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass Centipede St. Augustine Zoysiagrass	sulfosulfuron (0.035 - 0.059 lb)	Certainty (0.75 - 1.25 oz)	Nutsedge and kyllinga species <i>Poa annua</i>	A second application of 0.75 to 1.25 oz./acre may be made 4 to 10 weeks after initial treatment, if needed. Some chlorosis or stunting of the desirable turf may occur following application. Use of a nonionic surfactant is required. Certainty may be used for control of <i>Poa annua</i> prior to overseeding of perennial ryegrass.
	thiencarbazone+ iodosulfuron + dicamba	Celsius WG (2.5 - 4.9 oz)	Annual and perennial broadleaf weeds (see label for complete list).	For use on residential lawns, commercial lawns, golf courses, sports fields, parks, campsites, recreational areas, roadsides, school grounds, cemeteries, and sod farms.
	sulfentrazone + metsulfuron-methyl	Blindside 66 WDG (6.5 - 10 oz)	Broadleaf Weeds (see label for weeds controlled)	Labeled for use on residential and commercial lawns, athletic fields, sod farms and golf course fairways and roughs.
Centipede St. Augustine Seashore paspalum Zoysiagrass Bermudagrass (common and hybrid) Bahia	carfentrazone-ethyl (0.008 - 0.031 lb)	QuickSilver (0.9 - 2.1 fl oz) Aim (sod production)	Annual and perennial broadleaf weeds (see label for complete list).	Use as a stand-alone product for control of seedling broadleaf annual weeds or with other pre- and postemergence herbicides for control of annual and perennial broadleaf weeds. Some temporary discoloration may occur 3 to 7 days after application. Rapidly absorbed by weeds with symptoms often visible within 24 hours of application.
Bermudagrass Centipede Seashore paspalum Zoysiagrass	sulfentrazone + quinclorac	Solitare (16 - 32 oz)	Annual and perennial broadleaf weeds and sedges (see label for complete list).	Selective weed control in turf including residential, commercial and industrial lawns, athletic fields, commercial sod farms, golf course fairways and roughs, and other non-crop areas.
	carfentrazone + quinclorac	SquareOne (12 - 18 oz)	Annual and perennial broadleaf weeds and sedges (see label for complete list).	Can be applied to newly seeded, sodded, or sprigged turfgrass at 7 days or more after emergence. Application to zoysiagrass should be delayed at least 14 days after emergence.
Bahia Bermudagrass Centipede Zoysiagrass	sulfentrazone+2,4-D +MCP+dicamba	Surge (2.75 - 3.25 pts)	Broadleaves (see label for complete list).	Not for use on St. Augustine. For centipede, use lower rates within specified range until turfgrass tolerance to injury can be determined.
	sulfentrazone + imazethapyr	Dismiss South 4SC (9.5 - 14.4 oz)	Control similar to that provided by Dismiss with added nutsedge control.	Labeled for use on residential and commercial lawns, athletic fields, sod farms, golf course fairways and roughs.

GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass Centipedegrass Zoysiagrass	flazasulfuron	Katana 25WDG (1.5 - 3.0 oz)	Broadleaves Sedges Some grass species including removal of winter overseeded cool season grasses.	Katana can be applied to turfgrass on golf course fairways and roughs, industrial parks, sod farms, cemeteries, professionally managed sports fields, and non-residential commercial lawns.

Postemergence Herbicide Efficacy Ratings																					
E = excellent (>89%) control; F = Fair to good (70 to 89%), good control sometimes with high rates, however a repeat treatment 1 to 3 weeks later each at the standard or reduced rate is usually more effective; P = poor (<70%) control in most cases. Not all weeds have been tested for susceptibility to each herbicide listed. A = annual, B = biennial; P = perennial; SA = summer annual; WA = winter annual.																					
	Lifecycle	atrazine	bentazon	bispyribac-sodium	bromoxynil	carfentrazone	carfentrazone + quinclorac	clethodim	clopyralid	2,4-D	2,4-D + dicamba	dicamba	diclofop	ethofumesate	fenoxaprop	flazasulfuron	fluazifop	fluroxypyr	foramsulfuron	glufosinate	glyphosate
GRASS and "GRASS-LIKE" WEEDS																					
bahiagrass	P	P	P		P	P		F	P	P	P	P	P	F	P		F	P	P	P	G
bermudagrass	P	P	P		P	P		F-G	P	P	P	P	P		P		G	P	P	P	E
dallisgrass	P	P	P		P	P		F	P	P	P	P	P		P			P	P	P	E
nutsedge, purple	P	P	P		P	P		P	P	F	P	P	P		P		P	P	P	P	G
nutsedge, yellow	P	P	G		P	P		P	P	F	P	P	P		P		P	P	P	P	E
torpedograss	P					P			P								P	P	P	F	
wild garlic/onion	P	P	P		P	P		P	P	G	G	F	P		P		P	P	P	P	G
ANNUAL GRASSES																					
<i>Poa annua</i>	WA	E	P	G	P	P		F	P	P	P	P	P	G	P		P	P	E	E	E
crabgrass	SA	F	P		P	P		E	P	P	P	P	P	F	G-E		G	P	P	E	E
crowfootgrass	SA	P	P		P	P		E	P	P	P	P		G-E			G	P	P	E	E
goosegrass	SA	P	P		P	P		F-G	P	P	P	P	E		G		G	P	G	E	E
sandbur	SA	P	P		P	P		G	P	P	P	P			G-E		G	P	P	E	E
BROADLEAF WEEDS																					
bittercress, hairy	WA							P	F		E	E	P		P		P		P		
black medic	A					G		P	E	E	P	E	P		P		P	F	P		
buttercups	WA,B,P	F	P		P	G		P	F-P	E-F	E-F	E-F	P		P		P		P		
carpetweed	SA	E						P		E	E	E	P		P		P		P		
carrot, wild	A,B							P	F	F	E-F	E	P		P		P		P		
catsear dandelion	P							P	E	E-F	E-F	E	P		P		P	F	P		

Postemergence Herbicide Efficacy Ratings

E = excellent (>89%) control; F = Fair to good (70 to 89%), good control sometimes with high rates, however a repeat treatment 1 to 3 weeks later each at the standard or reduced rate is usually more effective; P = poor (<70%) control in most cases. Not all weeds have been tested for susceptibility to each herbicide listed. A = annual, B = biennial; P = perennial; SA = summer annual; WA = winter annual.

	Lifecycle	atrazine	bentazon	bispyribac-sodium	bromoxynil	carfentrazone	carfentrazone + quinclorac	clethodim	clopyralid	2,4-D	2,4-D + dicamba	dicamba	diclofop	ethofumesate	fenoxaprop	flazasulfuron	fluzifop	fluroxypyr	foramsulfuron	glufosinate	glyphosate
chamberbitter (niruri)	SA,P	G	P					P	P	P	P-F	P-F	P		P		P		P		E
common chickweed	WA	E	G		P			P	E-F	P	G	E	P		P		P	F	P	G	E
corn speedwell	WA	E	P		G			P	P	F	F	F	P		P		P	P	P		E
cudweed		G			G	F		P	E	G-E	E	E	P		P		P	F	P		G
dandelion	P	F	P		P			P	F	E	G	E	P		P		P	F	P		E
dayflower, spreading	SA	G-E	G					P		F	F	F	P		P		P		P		
dichondra	P	E	P		P	F		P		G	G	G	P		P		P		P		E
docks	P	G	P			G		P	E	F	G	E	P		P		P		P	G	E
doveweed	SA	G-E	P		P			P		P	F	P	P		P		P	P	P		G
Florida betony	P	F-G	P		P	P		P		F	G	G	P		P		P	F	P		E
geranium, carolina	WA	E				G		P	P	E	E	E	P		P		P	P	P		
ground ivy	P		P		P			P		P-F	F	G	P		P		P	P	P		G
hawkweed	P					G		P		E-F	E-F	E-F	P		P		P		P		
henbit	WA	E	P		G	G		P		P	G	E	P		P		P	G	G		E
hop clovers	WA	E			F	G		P	E	F	G	E	P		P		P	G	P	G	E
knawel	WA							P		P	E-F	E	P		P		P		P		
knotweed	SA	E			F			P	E	P	G	E	P		P		P		P		E
lespedeza	SA	E						P	P	P-F	G	E	P		P		P	F	P		E
mallow	P	P	P					P		F	F-G	E	P		P		P		P		
mock strawberry	P	P	P					P		P	G	G	P		P		P		P	G	
mouseear chickweed	WA,P	G	P			G		P	F	P-F	G	E	P		P		P	F	P	G	E
mugwort	P	P	P					P		F	F	G	P		P		P		P		G
mustards	WA	E	G		G			P	P	E	G	E	P		P		P		P		E
parsley piert	WA	E	G		G			P		P		E	P		P		P	F	P		E
pearlwort	WA							P	P	E-F			P		P		P		P		
pennywort (dollarweed)	P	E	P		P	P		P	F	G	G	E	P		P		P	F	P		E
pepperweed, VA	WA	E			F	G		P		E	E	E	P		P		P	P	P		

Postemergence Herbicide Efficacy Ratings

E = excellent (>89%) control; F = Fair to good (70 to 89%), good control sometimes with high rates, however a repeat treatment 1 to 3 weeks later each at the standard or reduced rate is usually more effective; P = poor (<70%) control in most cases. Not all weeds have been tested for susceptibility to each herbicide listed. A = annual, B = biennial; P = perennial; SA = summer annual; WA = winter annual.

	Lifecycle	atrazine	bentazon	bispyribac-sodium	bromoxynil	carfentrazone	carfentrazone + quinclorac	clethodim	clopyralid	2,4-D	2,4-D + dicamba	dicamba	diclofop	ethofumesate	fenoxaprop	flazasulfuron	fluzifop	fluroxypyr	foramsulfuron	glufosinate	glyphosate
pigweed	WA	G	P		F-G	G		P		E	E	E	P		P		P		P	G	
plantains	P	F	P		P			P	F	E	G	F	P		P		P		P		E
shepherdspurse	WA		G		G			P	F	E	E	E	P		P		P		P		
spurges	SA	E	P		F	G		P	F	F	G	G	P		P		P	F	P		E
spurweed (burweed)	WA	E	E		G			P	E	G	G	E	P		P		P	F	P		E
thistles	B,P	P						P	E	E-F	E-F	E-F	P		P		P		P		
VA buttonweed	P	P	P		P	G		P	F-P	P	F	F	P		P		P	F	P		G
violets	P	P	P			G		P		P	F	F	P		P		P	P	P		
white clover	P	E	P			G		P	E	F	G	E	P	F	P		P	E	P	G	F
yellow woodsorrel (<i>Oxalis</i>)	P	G	P		F	G		P	F-P	P	F	G	P		P		P	F	P	G	E

Postemergence Herbicide Efficacy Ratings																						
E = excellent (>89%) control; F = Fair to good (70 to 89%), good control sometimes with high rates, however a repeat treatment 1 to 3 weeks later each at the standard or reduced rate is usually more effective; P = poor (<70%) control in most cases. Not all weeds have been tested for susceptibility to each herbicide listed. A = annual, B = biennial; P = perennial; SA = summer annual; WA = winter annual.																						
	Lifecycle	hexazinone	imazaquin	mesotrione	metribuzin	metsulfuron	pronamide	quinclorac	rimsulfuron	sethoxydim	simazine	sulfentrazone	sulfentrazone+imazethapyr	sulfentrazone+metsulfuron	sulfentrazone + proflaminate	sulfentrazone + quinclorac	sulfentrazone+2,4-D+MCPA+dicamba	sulfometuron	sulfosulfuron	thiencarbazone+iodosulfuron+dicamba	triclopyr + clopyralid	trifloxysulfuron
GRASS and "GRASS-LIKE" WEEDS																						
bahiagrass	P	P	P		P	E	P	P		F	P					P	P		F	P	P	F
bermudagrass	P	P	P		P	P	P	P		F	P					P	P		P	P	P	P
dallisgrass	P		P		P	P	P	F		P	P					P-F	P			F	P	F
nutsedge, purple	P		G		P	P	P	P		P	P	G	G			F	P	E	E	P	P	E
nutsedge, yellow	P		F-G	G	P	P	P	P		P	P	G	G		G	F	P	E	E	P	P	E
torpedograss	P							E		P	P					F	P			P	P	G
wild garlic/onion	P		E		P	E	P			P	P					F	P			G	P	G
ANNUAL GRASSES																						
<i>Poa annua</i>	WA		P-F		E	P	E	P	G	P	E					P	P	G	G	P	P	G
crabgrass	SA		P	E	F	P	P	E		E	P		G			F	P			F	P	F
crowfootgrass	SA		P		G	P	P			F-G	P					P	P			P	P	
goosegrass	SA		P	G	G	P	P	P		G	P	G	G		G	F	P			P	P	
sandbur	SA		F		G	P	P			G	P					P	P			F	P	
BROADLEAF WEEDS																						
bittercress, hairy	WA				G	E				P						F						E
black medic	A							E		P		G	G			F	G	G			F	E
buttercups	WA,B,P		G	G		E				P	F	G	G			F	F	G		G		E
carpetweed	SA			G		P				P	G	G	G			F	F	G			G	G
carrot, wild	A,B			E		E				P											G	
catsear dandelion	P			E						P						G					G	E

Postemergence Herbicide Efficacy Ratings

E = excellent (>89%) control; F = Fair to good (70 to 89%), good control sometimes with high rates, however a repeat treatment 1 to 3 weeks later each at the standard or reduced rate is usually more effective; P = poor (<70%) control in most cases. Not all weeds have been tested for susceptibility to each herbicide listed. A = annual, B = biennial; P = perennial; SA = summer annual; WA = winter annual.

	Lifecycle	hexazinone	imazaquin	mesotrione	metribuzin	metsulfuron	pronamide	quinclorac	rimsulfuron	sethoxydim	simazine	sulfentrazone	sulfentrazone+imazethapyr	sulfentrazone+metsulfuron	sulfentrazone + prodiamine	sulfentrazone + quinclorac	sulfentrazone+2,4-D+MCPP+dicamba	sulfometuron	sulfosulfuron	thiencarbazone+iodosulfuron+dicamba	triclopyr + clopyralid	trifloxysulfuron	
chamberbitter (niruri)	SA,P		P							P	F										G		
common chickweed	WA		E	E	G	E	G		F	P	E	G	G		F	F-G	G	G			G	E	E
corn speedwell	WA		P		E	E	G			P	E											P	E
cudweed			F							P	G	G	G		F	F					G	G-E	
dandelion	P			E		E	P			P	P	G	G		F	P	G				G	E-F	G
dayflower, spreading	SA		G			P-F		P		P	G		G				G						F-G
dichondra	P						P			P	P										G	E	G
docks	P			G		E	P			P	P	G	G		F	F	G		F			E	
doveweed	SA				F	P	P			P	F								G			P	
Florida betony	P						P			P	P						F		F			G	
geranium, carolina	WA		G			F-G				P	G	G	G		G				F-G			F	G
ground ivy	P			G			P			P	P	G	G		G		G		F-G			E-F	G
hawkweed	P									P							G					E	
henbit	WA		G	G	G	G	P		F	P	E	G	G		F	F	G		G		G	E	G
hop clovers	WA			G	G	P-F				P	E				F	G					G	E	G
knawel	WA		G							P		G	G		G								
knotweed	SA				G					P	G	G	G		F		G					E	
lespedeza	SA	G			E	E				P	G	G	G		F		G					E	
mallow	P						P			P	P	G	G		F		G					E	
mock strawberry	P						P			P	P												
mouseear chickweed	WA,P		G	E	E	E	P			P	P	G	G		F	P-F	G		G		G	E	E
mugwort	P						P			P	P												
mustards	WA				F	F	P			P	G		G						G		G		

Postemergence Herbicide Efficacy Ratings

E = excellent (>89%) control; F = Fair to good (70 to 89%), good control sometimes with high rates, however a repeat treatment 1 to 3 weeks later each at the standard or reduced rate is usually more effective; P = poor (<70%) control in most cases. Not all weeds have been tested for susceptibility to each herbicide listed. A = annual, B = biennial; P = perennial; SA = summer annual; WA = winter annual.

	Lifecycle	hexazinone	imazaquin	mesotrione	metribuzin	met sulfuron	pronamide	quinclorac	rimsulfuron	sethoxydim	simazine	sulfentrazone	sulfentrazone+imazethapyr	sulfentrazone+met sulfuron	sulfentrazone + prodiamine	sulfentrazone + quinclorac	sulfentrazone+2,4-D+MCP+dicamba	sulfometuron	sulfosulfuron	thiencarbazone+iodosulfuron+dicamba	triclopyr + clopyralid	trifloxysulfuron	
parsley piert	WA		G		E		P			P	E	G			G						G	E	
pearlwort	WA									P													
pennywort (dollarweed)	P					G	P	E	F	P	P					G	G			F-G	E	G	
pepperweed, VA	WA	G								P	G						G						
pigweed	WA	G		E		E-G				P	F	G	G		G		G				G		
plantains	P			G		G	P			P	P	G			G		G				G	E	
shepherdspurse	WA					G				P		G			G		G	G	G	G	E-F	G	
spurges	SA				E	E			F	P	G	G			G	G	G			F-G	E-F	G	
spurweed (burweed)	WA		E		G	E	P			P	E	F			F						G	E	E
thistles	B,P		G	G		F				P	P						G					E	
VA buttonweed	P						P			P	P						F				F-G	F	G
violets	P						P			P	P	G					F					F-G	E
white clover	P		F	F	F	E	P			P	P					G	F		G		H	E	G
yellow woodsorrel (<i>Oxalis</i>)	P			G		F-G	P			P	P	G			F	G					F-G	F-G	G

SEDGES	
Leaves composed of a blade, a sheath, and a ligule. Leaf sheath is closed. Ligule is often absent, when present is tiny. Stem is often triangular.	
Yellow Nutsedge, <i>Cyperus esculentus</i>	Perennial; underground runners; runners stop at tubers; tubers sweet to taste or with little flavor; seedhead yellow. Leaf tip needle-shaped.
Purple Nutsedge, <i>Cyperus rotundus</i>	Perennial; underground runners that continue from tuber to tuber, forming chains; tubers bitter to taste; seedhead purple or reddish. Leaf tip boat-shaped.
Globe Sedge, <i>Cyperus globulosus</i>	Perennial; forming clump; shiny green leaves; base hard; seedheads like small globes. Cylindric Sedge just as common; has all the same characteristics as the above differing only in the seedhead which is shaped like a cylinder with smaller and finer flowers.
Perennial (Green) Kyllinga <i>Cyperus brevifolius</i>	Mat forming perennial from reddish purple rhizomes. Leaves and stems, dark green. Seedhead simple, nearly round or oblong, with three short leaves just below. Reproduces by seed and rhizomes. Found in low areas or where moisture is excessive.
Annual Sedge <i>Cyperus compressus</i>	Annual; spreads by seed formed from clusters of flat spikes. Spikes greenish, sometimes glossy, up to 1-inch long.

SEDGE CONTROL				
GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass Zoysiagrass	trifloxysulfuron (0.00469 - 0.02 lb)	Monument 75WG (0.1 - 0.53 oz)	Purple and Yellow Nutsedge Green Kyllinga	0.33 to 0.53 oz/Ac for purple and yellow nutsedge and Kyllinga. Buffer water to pH 7 or less. Weed death may take 2 to 4 weeks or more under cool conditions. Lightly irrigate 4 hours or more after application to minimize tracking/movement if applications are made next to non-labeled cool-season turfgrass species. Multiple applications needed for hard-to-control weeds.
Centipedegrass St. Augustinegrass	mesotrione (0.125 - 0.25 lb)	Tenacity 4L Centipedegrass (5 - 8 fl oz) St. Augustinegrass (4 fl oz)	Yellow nutsedge	FOR SOD PRODUCTION USE ONLY. Efficacious when applied to young yellow nutsedge. Addition of a non-ionic surfactant required. Can be applied prior to seeding (one day before) centipedegrass or after cutting St. Augustinegrass. Some discoloration of St. Augustinegrass may occur.
Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	halosulfuron (0.03 - 0.06 lb)	SedgeHammer (0.67 - 1.33 oz)	Most nutsedges	Note the low use rate. Nutsedges should be actively growing when treated. Repeat application(s) 3 to 4 weeks apart will be needed for complete control. Now labeled for 4 applications per year with a maximum use rate of 5.33 oz product/acre/year.
	imazaquin (0.375 - 0.50 lb)	Image 70DG (8.6 - 11.4 oz)	Purple nutsedge, sandspur, wild garlic, some broadleaves	Do not apply to newly seeded, sodded, sprigged areas. Apply only to actively growing turfgrass. Not labeled for use on Bahiagrass or golf greens. Repeat applications may be required as weeds mature. Treated turf may have a compacted growth habit and seedhead formation may be inhibited. Do not use on overseeded turf.

SEDGE CONTROL				
GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	sulfosulfuron (0.035 - 0.059 lb)	Certainty (0.75 - 1.25 oz)	Nutsedge and Kyllinga	A second application of 0.75 to 1.25 oz/acre may be made 4 to 10 weeks after initial treatment, if needed. Some chlorosis or stunting of the desirable turf may occur following application. Use of a nonionic surfactant is required.
Bahiagrass Bermudagrass Centipedegrass Zoysiagrass	sulfentrazone + imazethapyr (0.29 - 0.45 lb)	Dismiss South (9.5 - 14.4 fl oz)	Nutsedge and broadleaves	May be used on seeded, sodded, or sprigged turfgrasses that are well established. May cause some temporary discoloration.
Bahiagrass Bermudagrass Centipedegrass Seashore paspalum St. Augustinegrass Zoysiagrass Perennial Ryegrass	sulfentrazone (0.25 - 0.375 lb)	Dismiss (8.0 - 12.0 fl oz) Spartan (sod)	Yellow Nutsedge Purple Nutsedge Kyllinga sps. Goosegrass Selected broadleaves	May cause temporary discoloration in bahiagrass, St. Augustinegrass, and zoysiagrass. Do not apply to weakened turfgrass.
Bahiagrass Bermudagrass Centipedegrass Seashore paspalum St. Augustinegrass Zoysiagrass	sulfentrazone + prodiamine (See comments)	Echelon (See comments)	Preemergence control of yellow nutsedge, kyllinga, annual sedge	Bahiagrass, centipedegrass, seashore paspalum, and zoysiagrass rate: 0.57 - 0.75 lb ai/A, 18-24 fl oz/A per application not to exceed 1.125 lb ai/A, 36 fl oz/A per calendar year. St. Augustinegrass rate: 0.57 lb ai/A (18 fl oz/A) followed by 0.57 lb ai/A (18 fl oz/A) 45 to 60 days later. Temporary discoloration may occur. Bermudagrass rate: 0.75 - 1.125 lb ai/A, 24 - 36 fl oz/A.
Bahiagrass Bermudagrass Centipedegrass St. Augustinegrass Zoysiagrass	S-metolachlor (1.1 - 2.2 lbs)	Pennant Magnum 7.62L (1.3 - 2.6 pts)	Preemergence control of yellow nutsedge and annual sedge	For use on golf course fairways, sod farms, and commercial lawns. The higher rate will be necessary for turf grown on high organic (i.e., muck) soils. For commercial St. Augustinegrass sod production, do not use more than once every 6 weeks and do not apply more than 4.2 pts./A/yr. Tank mixing with atrazine will increase the weed control spectrum. Do not use Pennant Magnum on golf greens, tees, or aprons or within 4 months of overseeding. Irrigate within 7 days after application.
	bentazon (1-2 lbs)	Basagran T&O 4L (1.5 - 2.0 pts)	Yellow nutsedge, globe sedge, annual sedge and some broadleaf weeds	Apply when yellow nutsedge is actively growing under good soil moisture conditions. Repeat applications will probably be necessary. Will not satisfactory control purple nutsedge. Not labeled for golf greens.
Seashore paspalum	halosulfuron (0.03 to 0.06 lb)	Sedgehammer 75WG (0.67 - 1.33 oz) Sanda 75WG (Sod production)	Most nutsedges	Nutsedge should be actively growing when treated. A second treatment, if needed, may be made 6 - 10 weeks after the initial treatment. Use a non-ionic surfactant.

SEDGE CONTROL				
GRASS	COMMON NAME (lbs ai/acre)	TRADE NAME (product rate/acre)	WEEDS CONTROLLED	COMMENTS
Bermudagrass Centipedegrass Seashore paspalum Zoysiagrass	sulfentrazone + quinclorac	Solitare (16 - 32 oz)	Annual and perennial sedges (see label for complete list).	Selective weed control in turf including residential, commercial and industrial lawns, athletic fields, commercial sod farms, golf course fairways and roughs, and other non-crop areas.
	carfentrazone + quinclorac	SquareOne 70 WDG (12 - 18 oz)	Grass and broadleaf weeds.	SquareOne is labeled for use on residential and commercial lawns, athletic fields, sod farms and golf course fairways and roughs.
	flazasulfuron	Katana (1.5 - 3.0 oz)	Most sedges	The use of a non-ionic surfactant at 0.25 percent by volume (1 qt/100 gal) provides maximum performance.

Sedge control and turf tolerance to various herbicides (Refer to Herbicide Label for Specific Species Listing).

Herbicide(s)*	Nutsedge Control				Turf Tolerance							
	Purple	Yellow	Annual	Kyllinga	Bermuda	Carpetgrass	St.Augustine	Bahiagrass	Centipede	Zoysia	Seashore Paspalum	Overseed Ryegrass/ Blends
Basagran T/O	P	G	G	F-G	S	S	S	S	S	S	NR	S-I
Certainty	E	E	E	E	S	NR	S	NR	S	S	NR	D
Dismiss/Spartan	G	E	E	E	S	S	I	S	S	S	S	S
Dismiss South	G	E	E	E	S	NR	NR	I	S	S	NR	NR
Echelon	G	E	E	E	S	S	I	S	S	S	S	S
Image	G	G	G	G	I	I	S	D	S	S	NR	D
Katana	G	G	G	G	S	NR	NR	NR	S	S	S	D
Sedgehammer/Sandea	G	G	G	G	S	--	S	S	S	S	S	--
Solitare					S	NR	NR	NR	S	S	S	S
Monument	E	E	E	E	S	NR	NR	NR	NR	S	NR	D
Tenacity	NR	G	NR	NR	SD	NR	I	D	S	D	D	S
Trimec Plus	P	G	G	F	S-I	D	D	D	D	S-I	NR	D

S=Safe at labeled rates; I=Intermediate safety, use at reduced rates; SD=Safe if fully dormant; D=Damaging, do not use; NR=Not Registered for use on this turfgrass. G=good; F=fair; P=poor.

*Repeat applications are necessary for complete control from all herbicides.

Turfgrass Tolerance to Postemergence Herbicides (Refer to Herbicide Label for Specific Species Listing) ¹								
Herbicide	Bahiagrass	Bermudagrass	Carpetgrass ²	Centipede	Seashore Paspalum	St. Augustine	Zoysiagrass	Overseed Rye/Blends
atrazine (Aatrex)	NR	NR	I	S-I	NR	S-I	I	D
bentazon (Basagran)	S	S	S	S	NR	S	S	S-I
bispyrabac-sodium (Velocity)	NR	S ³	NR	NR	NR	NR	NR	S
bromoxynil (Buctril)	S	S	S	S	NR	S	S	S
carfentrazone (Quicksilver)	S	S	NR	S	S	S	S	S
carfentrazone+quinclorac (SquareOne)	NR	S	NR	S	S	NR	S	NR
clopyralid (Lontrel)	S	S	NR	S	S	S	S	S
2,4-D	S	S	I	I	NR	I	S	S-I
2,4-D+dicamba	S	S	I	I	NR	I	S	S-I
2,4-D+2,4-DP	S	S	I	I	NR	I	S	I-D
2,4-D+MCP	S	S	I	I	NR	I	S	I-D
2,4-D+MCP+dicamba	S	S	I	I	NR	I	S	I-D
2,4-D+MCP+2,4-DP	S	S	I	I	NR	I	S	I-D
2,4-D+MCP+dicamba+MSMA	D	S-I	D	D	NR	D	S-I	D
2,4-D+clopyralid+dicamba	S	S	I	I	NR	S-I	S	S
dicamba (Banvel)	S	S	I	I	NR	I	S	I
diclofop (Illoxan)	NR	S	NR	NR	NR	NR	NR	NR
ethofumesate (Prograss)	NR	D	--	NR	NR	I	NR	D
fenoxaprop (Acclaim)	I-D	I-D	D	D	NR	D	I	I
flazasulfuron (Katana)	NR	S	NR	S	NR	NR	S	D
fluazifop (Fusilade)	NR	NR	NR	NR	NR	NR	S	NR
fluroxypyr (Spotlight)	S	I	NR	S	NR	D	S	S
foramsulfuron (Revolver)	NR	S	NR	D	NR	NR	S	D
iodosulfuron+dicamba (Celsius WG)	NR	S	NR	S	NR	S	S	D
imazaquin (Image)	NR	S	NR	S	NR	S	S	NR
hexazinone (Velpar)	I	I-S	NR	NR	NR	NR	NR	D
MCPA+MCP+2,4-DP	S	S	I	I	NR	I	I	I-D

Turfgrass Tolerance to Postemergence Herbicides (Refer to Herbicide Label for Specific Species Listing) ¹								
Herbicide	Bahiagrass	Bermudagrass	Carpetgrass ²	Centipede	Seashore Paspalum	St. Augustine	Zoysiagrass	Overseed Rye/Blends
MCPP	S	S	I	I	NR	I	S	I
mesotrione (Tenacity)	NR	D	D	S	D	I	D	S
metribuzin (Sencor)	D	S-I	D	D	NR	D	NR	D
metsulfuron (Escort, Manor)	D	S	I	S	NR	S-I	S	D
penoxsulam (Lockup)	S	S	--	S	S	S	S	S
pronamide (Kerb)	NR	S	NR	NR	NR	NR	NR	D
pyraflufen ethyl (Octane)	NR	S	NR	S	NR	S	S	S
quinclorac (Drive)	D	I-S	--	D	S	D	S	S
rimsulfuron (TranXit)	NR	S	NR	NR	NR	NR	NR	NR
sethoxydim (Segment)	D	D	D	S	NR	D	D	D
simazine	NR	I	--	S	NR	S	S-I	D
sulfentrazone (Dismiss)	S	S	S	S	S	I	S	S
sulfentrazone+imazethapyr (Dismiss South)	I	S	NR	S	NR	NR	S	NR
sulfentrazone+metsulfuron-methyl (Blindside)	NR	S	NR	S	NR	S	S	NR
sulfentrazone+prodiamine (Echelon)	S	S	S	S	S	I	S	S
sulfentrazone+quinclorac (Solitaire)	NR	S	NR	S	S	NR	S	S
sulfentrazone+2,4-D+MCPP+dicamba	S	S	NR	I	NR	NR	S	S
sulfometuron (Oust)	I	I	NR	NR	NR	NR	NR	NR
sulfosulfuron (Certainty)	NR	S	NR	S	NR	S	S	D
thiencarbazon+iodosulfuron+dicamba (Celsius WG)	NR	S	NR	S	NR	S	S	D
triclopyr + clopyralid (Confront)	I	I	NR	I	NR	D	I	S
trifloxysulfuron (Monument)	NR	S	NR	NR	NR	NR	S	D

¹S=Safe at labeled rates; I=Intermediate safety, use at reduced rates; D=Damaging, do not use; NR=Not Registered for use on this turfgrass.

²Carpetgrass tolerance to herbicides listed has not fully been explored.

³Safe when applied to overseeded bermudagrass January 1 to April 15.

Presence of a herbicide in this listing does not constitute a recommendation. Trade names are used with the understanding that no endorsement is intended or no criticism is implied of similar products not mentioned. All chemicals should be used in accordance with the manufacturer's instructions.

Common and Trade Names of Turf Herbicides.

Common Name	Manufacturer / Distributor	Trade Name(s)
Atrazine	Syngenta, Ortho, Security, + others	Aatrex, Atrazine Plus, Purge II, + others
Benefin	Dow AgroSciences + others	Balan, Crabgrass Preventer, + others
Benefin & oryzalin	Dow AgroSciences	XL 2G
Benefin & oxadiazon	Regal Chemical	RegalStar
Benefin & trifluralin	Verdicon / UHS	Team 2G
Bensulide	Gowan, Scotts, + others	Betasan, Pre-San, Bensumec 4, Weedgrass Preventer, + others
Bentazon	BASF, LESCO	Basagran T&O, Lescogran 4L
Bispyrabac	Valent USA	Velocity
Bromoxynil	Bayer	Buctril 2L
Carfentrazone	FMC	Quicksilver/Aim
Carfentrazone + quinclorac	FMC	SquareOne
Carfentrazone + 2,4-D + MCPP + Dicamba	PBI/Gordon	SpeedZone, SpeedZone - Southern
Carfentrazone + MCPA + MCPP + Dicamba	PBI/Gordon	Power Zone
Clethodim	Valent USA Corp.	Envoy
Clopyralid	Dow AgroSciences	Lontrel T&O
2,4-D	Bayer, NuFarm, + others	Many
2,4-D+Clopyralid + Dicamba	NuFarm	Millennium Ultra 2
DCPA	Syngenta	Dacthal 75WP, Garden Weed Preventer, + others
Dicamba	Syngenta, Scotts, + others	Vanquish 4L, K-O-G Weed Control, + others
Diclofop	Bayer	Illoxan 3EC
Dimethenamid-P	BASF	Tower
Dithiopyr	Dow AgroSciences, Quali-Pro	Dimension Ultra, QP Dithiopyr
Diquat	Syngenta	Reward Landscape & Aquatic Herbicide
Ethofumesate	Bayer	Prograss 1.5L
Fenoxaprop	Bayer	Acclaim

Common Name	Manufacturer / Distributor	Trade Name(s)
Flazasulfuron	PBI/Gordon	Katana
Fluazifop	Syngenta	Fusilade II
Fluroxypyr	Dow AgroSciences	Spotlight
Foramsulfuron	Bayer	Revolver
Glufosinate	Bayer	Finale 1L
Glyphosate	Monsanto, Syngenta, + others	RoundUp, Touchdown PRO, Glyphomaxx
Halosulfuron	Gowan	Sedgehammer 75DG / Sandea 75DG
Hexazinone	DuPont	Velpar 2L
Imazaquin	BASF	Image 70DG
Indaziflam	Bayer	Specticle 20WSP
Isoxaben	Dow AgroSciences	Gallery 75DF
MCPP	PBI/Gordon, Ortho, + others	Mecomec 4
MCPP, 2,4-D + dicamba + MCPA and/or 2,4-D	PBI/Gordon, LESCO, NuFarm, Sierra,	Trimec Southern/3-Way Selective/Eliminate DG/33, others
Mesotrione	Syngenta	Tenacity 4L
Metribuzin	Bayer	Sencor 75DF
Metolachlor	Syngenta	Pennant Magnum
Metsulfuron	NuFarm, Quali-Pro	Manor, QP MSM
Napropamide	United Phosphorus, UHS	Devrinol 50DF, Ornamental Herbicide 2G
Oryzalin	UPI, Quali-Pro	Surflan AS, QP Oryzalin
Oxadiazon	Bayer, Quali-Pro	Ronstar 2G, 50WP, QP Oxadiazon 2G, 50 WP
Pendimethalin	LESCO, BASF, Scotts	Pre-M, Pendulum, Pendulum AquaCap, Southern Weedgrass
Penoxulam	Dow AgroSciences	Lockup
Prodiamine	Syngenta, NuFarm, Quali-Pro	Barricade 65WG, 4FL, ProClipse, QP Prodiamine
Pronamide	Dow AgroSciences	Kerb 50WP
Pyraflufen ethyl	SePRO	Octane 2% SC
Quinclorac	BASF	Drive

Common Name	Manufacturer / Distributor	Trade Name(s)
Quinclorac + dicamba + MCPP	BASF	Onetime Herbicide
Rimsulfuron	DuPont	TranXit
Sethoxydim	BASF	Segment
Simazine	Syngenta + others	Princep Liquid, others
Sulfentrazone	FMC	Dismiss
Sulfentrazone + 2,4-D + MCPP + Dicamba	PBI Gordon	Surge Broadleaf Herbicide
Sulfentrazone + imazethapyr	FMC	Dismiss South
Sulfentrazone + metsulfuron	FMC	BlindSide
Sulfentrazone + prodiamine	FMC	Echelon
Sulfentrazone + quinclorac	FMC	Solitare
Sulfometuron	DuPont	Oust
Sulfosulfuron	Monsanto	Certainty
thiencarbazone + Iodosulfuron-methyl-sodium + dicamba	Bayer	Celsius WG
Triclopyr + clopyralid	Dow AgroSciences	Confront
Trifloxysulfuron	Syngenta	Monument

Plant Growth Retardants for Fine Turf and Roadsides/Utilities

J. Bryan Unruh, Ph.D. and Barry J. Brecke, Ph.D.
Extension Turfgrass Specialist and Research Weed Scientist

Plant growth retardants (PGR's) or inhibitors are increasingly being used to suppress seedheads and leaf growth due to rising mowing costs and danger posed to operators and other personnel. Traditionally, plant growth retardants have been used in the South to suppress bahiagrass (*Paspalum notatum* Flugge.) seedhead production exclusively in low maintenance areas such as highway roadsides, airports, and golf course roughs. However, in recent years, new chemicals which may be used in higher maintained commercial situations have been developed.

Several undesirable characteristics which have been associated with growth retardants include: phytotoxicity (burn) of treated leaves from 4 to 6 weeks following applications; reduced recuperative potential from physical damage to treated turf; and increased weed pressure due to reduced competition from treated plants. Normally, growth retardants are used in low maintenance areas; therefore, these undesirable characteristics do not pose a problem to most managers. However, several growth regulatory materials have recently been developed for use on hybrid bermudagrass fairways and St. Augustinegrass. Vertical topgrowth (clippings) is suppressed, but horizontal spread (runners) is not. Therefore, turf recovery from golf club divots and other injuries occurs while topgrowth remains suppressed. Other uses involve areas where mowing has been discontinued due to heavy rains, equipment failure, etc., but topgrowth remains suppressed if the grass is treated. **Note: These retardants used on hybrid bermudagrass and St. Augustinegrass do not satisfactorily suppress seedhead development.**

PGRs are separated into two groups, Type I and Type II, based on their method of growth inhibition or suppression. Type I inhibitors are primarily absorbed through the foliage and inhibit cell division and differentiation in meristematic regions. They are inhibitors of vegetative growth and interfere with seedhead development. Their growth inhibition is rapid, occurring within 4 to 10 days, and lasts 3 to 4 weeks, depending on application rate. Mefluidide, chlorflurenol, and maleic hydrazide are examples of Type I inhibitors that inhibit mitosis in growth and development. Other Type I PGRs that inhibit plant growth and development through interruption of amino acid or organic acid biosynthesis are herbicides used at low rates. Being herbicides, their margin of safety is narrow and are very rate dependent. Examples of Type I herbicide regulators include glyphosate, imidazolinones, sulfonyleureas, sethoxydim, and fluzafop.

Type II inhibitors are generally root absorbed and suppress growth through interference of gibberellic acid bio-synthesis, a hormone responsible for cell elongation. Type II PGRs are slower in growth suppression response, but their duration is usually from 4 to 7 weeks, again, depending on application rate. Type II PGRs have little effect on seedhead development and results in miniature plants. Paclobutrazol and flurprimidol are root absorbed Type II PGRS while trinexapac-ethyl is a foliar absorbed Type II PGR and is systemically translocated to the site of activity. Fenarimol is a type II fungicide that also suppresses *Poa annua* on putting greens.

Root absorbed PGRs are activated by irrigation or rainfall after application and have less likelihood of over-lap leaf burn. Foliar absorbed materials (e.g., mefluidide, MH, and trinexapac-ethyl) require uniform and complete coverage for uniform response and must be leaf absorbed before irrigation or rainfall occurs. Usually low gallonage is used for foliar absorbed materials to minimize runoff from the leaf surface while high gallonage is used for root absorbed materials.

Timing of application for seedhead suppression is somewhat important. Applications made after seedhead emergence may not be effective. For bahiagrass, mow the area as seedheads initially emerge (usually in late May to early June) to knock these and weeds present down. Begin plant growth retardant treatment about two weeks following mowing or just prior to new seedhead appearance. Additional applications 6 to 8 weeks later may be required if new seedheads begin to emerge. A complete weed control program must accompany any plant growth retardant use. Typically, annual broadleaf weeds will become established in PGR use areas as the treated grass is not actively growing, therefore, is not providing its usual competition. Normally, 2,4-D and/or dicamba is included in this broadleaf weed control. Other postemergence herbicides such as Velpar, for grass weed control, may also be incorporated in low maintenance bahiagrass areas. The following tables list chemicals, application rates, and general remarks about each product used to suppress plant growth.

Chemical (Trade Names)	Active Ingredient per Acre	Amount of Product Per Acre	Remarks
Bermudagrass and St. Augustinegrass			
mefluidide (Embark 2S) (Embark Turf and Ornamental Growth Regulator)	0.125 - 1.0 0.125	½ to 4 pts/15-150 gal water 5 pts (St. Augustinegrass)	Foliar absorbed. Apply to common bermudagrass (4 pts/A Embark 2S) and St. Augustinegrass (Embark T&O) only. Apply in spring approximately 2 weeks before seedhead appearance. Do not apply to turf within 4 growing months after seeding, and do not reseed within 3 days after application. Treated turf may appear less dense and temporarily discolored. Adding 1 to 2 qts of a nonionic surfactant per 100 gal of spray solution may enhance suppression; however, discoloration may also be increased. <i>Poa annua</i> seedhead control in fairways is with ½ pt/A in early January. Iron applications may lessen discoloration. Read and follow label recommendations before use.
flurprimidol (Cutless 50 WP)	0.25 to 1.5 lbs	0.5 to 3.0 lb	Root absorbed. Apply to bermudagrass or zoysiagrass golf course fairways, hard-to-mow and trimmed areas. Provides 4 to 8 week suppression. Must be uniformly applied and irrigated in with ½ inch water. Flurprimidol does not completely control seedheads. Temporary turf discoloration may follow this treatment. St. Augustinegrass, bahiagrass, and common bermudagrass require the higher rate. Repeat applications every 4 weeks on Tifway bermudagrass with 0.5 lb/A will minimize turf injury. Good growth regulation and turf enhancement have been noted with half-rates of the Cutless + PrimoMAXX tank mix. Not recommended for bermudagrass golf greens.
trinexapac-ethyl (Primo MAXX and others)	0.1 to 0.75	9 to 88 oz / A	Foliar absorbed. Low rates are for hybrid bermudagrass, centipedegrass, and St. Augustinegrass; medium rates are for common bermuda while the high rate is for bahiagrass and for edging and banding of bermudagrass and St. Augustinegrass. One hour rain-free period is needed after application. Mowing one week after application improves results and appearance as will repeat applications in two to four weeks. Do not exceed 21 pints/A per year (WSB = 174 fl. oz/A/year). Seedhead suppression is provided only for hybrid bermudagrass. Temporary turf discoloration may follow treatment. Do not add a surfactant.
Trinexapac-ethyl + 5-0-10 fertilizer (Governor)	0.02 to 0.44 lbs	36 - 720 lbs / Ac	A granular formulation of trinexapac-ethyl for professional use, which could also be purchased for homeowner use. This warm-season-turf formulation works to reduce clippings and mowing frequency like the sprayable product.
Flurprimidol + Trinexapac-ethyl (Legacy)	Flurprimidol - 0.086 - 0.172 + Trinexapac-ethyl - 0.032 - 0.064	10 - 20 fl oz/A	Foliar and root absorbed. Apply to hybrid bermudagrass, zoysiagrass, and Seashore paspalum. Use lower rates for zoysiagrass and TifSport bermudagrass. Repeat applications every 3 to 6 weeks. Rainfall or irrigation should be delayed at least 2 hours after application or until product has dried on the leaf surface but should occur within 24 hours after application. Temporary turf discoloration may follow treatment. Do not apply to bermudagrass putting greens or overseeded bermudagrass putting greens.

Chemical (Trade Names)	Active Ingredient per Acre	Amount of Product Per Acre	Remarks
paclobutrazol (TGR Turf Enhancer 50WP)	0.5 - 1.0	1.0 - 1.5 lb in 43 - 100 gal water	Root absorbed. Apply to well-maintained St. Augustinegrass or hybrid bermudagrass fairways. Do not apply to bermudagrass greens. Do not apply to saturated soils and treat only dry foliage. Repeat applications 8 weeks apart may be made. Read and follow recommendations before use.
Turf Growth Regulator + fertilizer 0.82%	0.6 - 0.9 lb	73 to 110 lb	
Trimmit 2SC	0.5 - 0.75 lb	32 to 48 oz	
Foliar Suppression of Overseeded Bermudagrass			
trinexapac-ethyl (Primo MAXX and others)	0.1 - 0.75	0.75 - 6 pints in 20 to 100 gal water	Apply trinexapac-ethyl before verticutting, scalping, spiking, or other similar operations or 1 to 5 days before seeding. Trinexapac-ethyl will allow the use of less severe base preparation practices. Check the label for the type of turf and setting (greens, tees, electron transport chain.) Use upper- end label rates when there is strong bermudagrass competition. Use normal seeding rates and maintain fertility.
mefluidide (Embark 2S)	0.125	0.5 pts in 15-150 gal water	Root absorbed. Do not apply to turf within 4 growing months after seeding, and do not reseed within 3 days after application. Treated turf may appear less dense and temporarily discolored. Adding 1 to 2 qts of a nonionic surfactant per 100 gal of spray solution may enhance suppression; however, discoloration may also be increased. <i>Poa annua</i> seedhead control in fairways is with ½ pt/A in early January. Iron applications may lessen discoloration. Read and follow label recommendations before use.
flurprimidol (Cutless 50W)	0.375 - 1.50 lb	0.75 - 3.0 lbs in 50 to 200 gal water	Root absorbed. Apply to zoysiagrass or bermudagrass in late spring-early summer and, or late summer-early fall. Time the second application at least 3 months before expected dormancy. Do not apply to putting greens. Do not exceed 1½ lb/A per application on sandy soils. Irrigate with ½in. water & resume mowing 3 to 5 days after application.
Flurprimidol + Trinexapac-ethyl (Legacy)	Flurprimidol - 0.129 - 0.258 + Trinexapac-ethyl - 0.048 - 0.096	15 - 30 fl oz/A	Foliar and root absorbed. Apply to perennial ryegrass overseeded turf areas. Do NOT apply to overseeded bermudagrass golf greens. For bermudagrass fairways overseeded with perennial ryegrass, applications of Legacy should be delayed until perennial ryegrass is well established (3 to 4 weeks after germination). Final spring application should be made a minimum of 4 weeks prior to expected bermudagrass green-up. Use lower rates for initial applications. Repeat applications every 3 to 6 weeks. Suppression of annual bluegrass (<i>Poa annua</i>) populations will occur with programmed applications of Legacy. Rainfall or irrigation should be delayed at least 2 hours after application or until product has dried on the leaf surface but should occur within 24 hours after application. Temporary turf discoloration may follow treatment. Do not apply to bermudagrass putting greens or overseeded bermudagrass putting greens.

Chemical (Trade Names)	Active Ingredient per Acre	Amount of Product Per Acre	Remarks
paclobutrazol (TGR Turf Enhancer 50 WP)	0.25 lb	0.5 lb in 40 - 100 gal water	Root absorbed. Repeat applications may be made 3 weeks apart. Do not use if <i>Poa annua</i> exceeds 70%. Application should be in early January.
Trimmit 2SC	0.1 - 0.5 lbs	6.4 - 32 oz	Root absorbed. Repeat applications may be made 4 to 6 weeks apart. Do not apply less than 2 weeks before and 6 weeks after overseeding. Read label for all directions. Do not apply to actively growing bermudagrass greens.

Read and follow all label recommendations before use. Products listed are for use by professional turf managers only. Trade and brand names are used for information only. The Florida Cooperative Extension Service, IFAS, and the University of Florida do not guarantee nor warrant the standard of any product mentioned; neither do they imply approval of any product to the exclusion of others which may also be suitable.

ACTIVATED CHARCOAL FOR PESTICIDE DEACTIVATION

J. Bryan Unruh and Barry J. Brecke
Extension Turfgrass Specialist and Research Weed Scientist

Introduction

Occasionally an accidental spill will occur that must be cleaned up to continue growing grass. Or in a combination weed control and grass seeding program, it is necessary to stop the activity of an applied herbicide to successfully seed a grass. Charcoal is a very porous, soft, black substance made by heating, in a restricted amount of air, substances containing carbon. This is most often derived from hardwood trees and coconut shells. Charcoal adsorbs 100 to 200 times its own weight. The adsorption capacity is developed by activating the charcoal by heating. Its ultra fine-grained and irregular shape give charcoal an enormous surface areas.

Activated charcoal comes in handy for binding, thus, deactivating some herbicides. Activated charcoal will reduce the available level of most organic pesticides in the soil; however, it is considered ineffective for inorganic pesticides such as arsenates, lead compounds, sodium chlorate, sulfur, borax, etc., and water-soluble organic pesticides such as, but not limited to, aminotriazole, MSMA, and DSMA. In order to have effective application of activated charcoal, it is important to have the spraying equipment clean and in good operating condition. It is a good idea to keep a bag or two of activated charcoal in stock at all times when managing fine turf so it can be applied almost immediately instead of having to wait several days for delivery after an accidental spill or application.

General Mixing Instructions:

For application convenience, it is recommended that activated charcoal be applied as a water slurry. To minimize dusting, always add activated charcoal to water slowly, keeping the bag as close to the water surface as possible.

Spray Application

1. Make sure spray equipment, tubing, and nozzles are completely clean. Screens should be removed if practical.
2. The final spray mixture should contain 1 to 2 lbs of charcoal per gallon of water.
3. Add sufficient water to begin moderate agitation. Simultaneously add the balance of required water and charcoal. Continue agitation until a uniform mixture is obtained.
4. Maintain moderate agitation while spraying.

Application	Recommendation	Comments
Spills	For reducing the effects from spills of organic pesticides, some petroleum products, and hydraulic fluids.	Use 100 lbs of activated charcoal to every pound of active material spilled but no less than two pounds per 150 sq.ft. (600 lbs/acre) of contaminated area. If the active material has not been diluted with water at the time of spill, apply the charcoal directly as a dry power. If the active material has been diluted with water, apply the activated charcoal in a slurry with a sprinkle can or common sprayer equipment. The charcoal must be incorporated into the contaminated soil, preferably to a depth of six inches. With severe spills, some of the contaminated soils may need removed prior to application.
'Deactivating' turf herbicides and Soil warming	Turf areas that have been treated with preemergence herbicides can be reseeded earlier than normal by treating with activated charcoal.	Whenever it is desirable to terminate the effect of a preemergence crabgrass herbicide, apply charcoal slurry at a rate of one lb per gallon of water for each 150 sq.ft. Water the slurry into the soil. Make sure the grass is washed free of heavy charcoal deposits. Where possible, it is desirable to rake the charcoal into the soil thoroughly. The area can be seeded 24 hrs after treatment.

Powdered activated charcoal is available as:

'DARCO Gro-Safe'
Norit Americas., Inc.
3200 University Avenue
Marshall, TX 57670
1-800-641-9245
www.norit-americas.com

'52 Pickup'
Brandt Consolidated
2935 South Koke Mill Road
Springfield, IL 62711
1-271-547-5840
www.brandtconsolidated.com

EFFICACY OF HERBICIDES ACTIVE INGREDIENTS AGAINST AQUATIC WEEDS

K. Langeland, M. Netherland, W. Haller, and T. Koschnick

Ken Langeland, professor, Agronomy Department, Center for Aquatic and Invasive Plants; Michael Netherland, courtesy associate professor, Center for Aquatic and Invasive Plants; William Haller, professor, Center for Aquatic and Invasive Plants; Tyler Koschnick, research assistant professor, Agronomy Department; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.

Please refer to:

<http://edis.ifas.ufl.edu/AG262>

Pesticide Calibration Formulas and Information
J. Bryan Unruh
Extension Turfgrass Specialist

$$\text{Acres covered/hour:} = \text{MPH} \times \text{Swath (ft)} \times 0.1212$$

$$= \frac{\text{MPH} \times \text{Swath (ft)}}{8.25}$$

$$\text{Gallons Per Acre (GPA):} = \frac{\text{GPM} \times 495}{\text{MPH} \times \text{Swath (ft)}}$$

$$= \frac{\text{GPM per nozzle} \times 495}{\text{MPH} \times \text{nozzle spacing (ft)}}$$

$$= \frac{\text{GPM per nozzle} \times 5940}{\text{MPH} \times \text{nozzle spacing (inches)}}$$

$$= \frac{\text{fl.oz. collected per nozzle} \times 4084}{\text{ft. traveled} \times \text{nozzle spacing (inches)}}$$

$$= \frac{\text{fl.oz collected per nozzle in 100 ft} \times 40.8375}{\text{nozzle spacing (inches)}}$$

$$= \frac{\text{gallons per 1000 sq.ft.}}{0.023}$$

$$= \frac{\text{gallons collected per nozzle} \times \text{no. of nozzles} \times 43560}{\text{ft. traveled} \times \text{Swath (ft)}}$$

$$\text{Gallons per 1000 sq.ft.} = 0.023 \times \text{GPA}$$

$$\text{Ounces per 1000 sq.ft.} = 2.94 \times \text{GPA}$$

$$\text{Gallons Per Minute (GPM):} = \frac{\text{GPA} \times \text{MPH} \times \text{Swath (ft)}}{495}$$

$$= \frac{\text{GPA} \times \text{MPH} \times \text{nozzle spacing (inches)} \times \text{no. nozzles}}{5940}$$

$$= \frac{\text{fl.oz per minute}}{128}$$

$$\text{GPM/Nozzle:} = \frac{\text{GPA} \times \text{MPH} \times \text{nozzle spacing (inches)}}{5940}$$

$$= \frac{\text{GPA} \times \text{MPH} \times \text{nozzle spacing (ft)}}{495}$$

$$= \frac{\text{Test jar fl.oz} \times 0.46875}{\text{seconds to fill test jar}}$$

$$= \frac{7.5}{\text{seconds to fill 1 pint (16 fl.oz.)}}$$

$$= \frac{15}{\text{seconds to fill 1 quart (32 fl.oz.)}}$$

$$\text{Minutes/Acre:} = \frac{495}{\text{MPH} \times \text{Swath (ft)}}$$

$$\text{Minutes/load:} = \frac{\text{gallons/load} \times 495}{\text{MPH} \times \text{GPA} \times \text{Swath (ft)}}$$

$$\text{Travel Speed} = \frac{\text{Distance traveled (ft)} \times 0.68}{\text{time (seconds) to travel distance}}$$

(Miles Per Hour, MPH)

$$\text{Acres covered per tank:} = \frac{\text{Gallons per tank}}{\text{GPA}}$$

$$\text{Material needed per tank} = \frac{\text{rate/A} \times \text{gallons/tank}}{\text{GPA}}$$

Flow Rate (as influenced by pressure):

$$\frac{GPM_1}{GPM_2} = \frac{\sqrt{PSI_1}}{\sqrt{PSI_2}} \quad \text{or} \quad GPA_2 = GPA_1 \times \sqrt{\frac{PSI_2}{PSI_1}} \quad \text{or} \quad PSI_2 = PSI_1 \times \left(\frac{GPA_2}{GPA_1}\right)^2$$

For any change in travel speed (mph), calculate the resulting GPA₂ by:

$$GPA_2 = \frac{GPA_1 \times MPH_1}{MPH_2} \quad \text{or} \quad \frac{GPA_1}{GPA_2} = \frac{MPH_2}{MPH_1} \quad \text{or} \quad MPH_2 = \frac{GPA_1 \times MPH_1}{GPA_2}$$

Fluid Application:

lbs/acre nutrient applied	= 0.226464 x element concentration (ppm) x acre inches of solution applied
PPM	= $\frac{1,000,000 \times \text{lbs ai used}}{\text{gal/tank} \times 8.34}$ or $\frac{\text{wt. of material to be used (lbs)} \times 1,000,000}{\text{wt. of tank mixture (lbs)}}$
	= $\frac{1,000,000 \times \text{oz commercial material used} \times \% \text{ ai (decimal)}}{\text{gal/tank} \times 8.34 \times 16}$
	= $\frac{1,000,000 \times \text{fl.oz. used} \times \text{lb ai/gal}}{\text{gal/tank} \times 8.34 \times 128}$
lbs nutrients applied/acre	= ppm of the element in the water x acre-inches water applied x 0.226464
lb ai to use per tank	= $\frac{\text{PPM desired} \times \text{gal/tank} \times 8.34}{1,000,000}$ or $\frac{\text{ppm desired} \times \text{gal/tank} \times 8.34}{1,000,000 \times \% \text{ ai}}$
lb commercial material to use per tank	= $\frac{\text{PPM desired} \times \text{gal/tank} \times 8.34}{1,000,000 \times \% \text{ ai (decimal)}}$ or $\frac{\% \text{ desired} \times \text{gal/tank} \times 8.34}{\% \text{ ai (decimal)}}$
fl. oz. to use per tank	= $\frac{\text{PPM desired} \times \text{gal/tank} \times 8.34 \times 128}{1,000,000 \times \text{ai per gal}}$
gal commercial material to use per tank	= $\frac{\text{ai (decimal)} \times 8.34 \text{ gal/tank}}{\text{ai per gal} \times 100}$
% ai in a spray mix	= $\frac{\text{lbs. commercial material used} \times \% \text{ ai (decimal)}}{\text{gal/tank} \times 8.34}$
gal commercial material for total treated acres	= $\frac{\text{PPM desired} \times \text{GPA} \times \text{acres} \times 8.34}{1,000,000 \times \text{lb ai/gal}}$ Active Ingredients (ai)
lbs commercial material/acre	= $\frac{\text{lbs ai to be applied per acre}}{\% \text{ ai of material}}$
gal commercial material/acre	= $\frac{\text{lbs ai to be applied per acre}}{\text{lbs ai per gallon}}$
gal commercial material/tank	= $\frac{\text{gallons/tank} \times \text{lb ai to be applied per acre}}{\text{gallons/acre} \times \text{lbs ai per gallon}}$

Time (seconds) required to cover a specific distance to obtain a desired speed (MPH).

Desired MPH	Feet per minute	Time Required (Seconds) to Travel a Distance of		
		100 ft.	200 ft.	300 ft.
2.0	176	34	68	102
2.5	220	27	54	81
3.0	264	23	45	68
3.5	308	20	39	58
4.0	352	17	43	51
4.5	395	15	30	45
5.0	440	14	27	41
6.0	528	--	23	34
7.0	616	--	19	29
8.0	704	--	17	26
9.0	792	--	15	23

Metric Prefix Definitions

tera	=	10 ¹²	deci	=	10 ⁻¹
giga	=	10 ⁹	centi	=	10 ⁻²
mega	=	10 ⁶	milli	=	10 ⁻³
kilo	=	10 ³	micro	=	10 ⁻⁶
hecto	=	10 ²	nano	=	10 ⁻⁹
deca	=	10 ¹	pico	=	10 ⁻¹²
basic metric unit = 1					

Approximate Rates of Application Equivalents


	<u>weights</u>	
1 oz/ft ²	= 2722.5 lbs/A	
1 oz/yd ²	= 302.5 lbs/A	
1 oz/100 ft ²	= 27.2 lbs/A	
1 oz/1000 ft ²	= 43.46 oz/A	= 2.72 lbs/A
1 lb/A	= 1 oz/2733 ft ²	= 8.5 g/1000 ft ²
100 lb/A	= 2.5 lb/1000 ft ²	
1 yd ³ sand	≈ 1.3 to 1.5 tons	
1 bushel	= 1¼ ft ³ = 0.046 yd ³	

	<u>Liquid</u>
1 oz/1000 ft ²	= 43.56 oz/A = 1.4 qt/A
1 pt/1000 ft ²	= 5.4 gal/A
100 gal/A	= 2.3 gal/1000 ft ² = 1 qt/100 ft ²


Approximate Weight of Dry Soil

Type	lbs/ft ³	lbs/acre (6 inches deep)
sand	100	2,143,000
loam	80-95	1,714,000
clay or silt	65-80	1,286,000
muck	40	860,000
peat	20	430,000


			<u>Helpful Calculations and Formulas:</u>	
Rectangle, square or parallelogram:	area	=	length (L) x width (W)	
Trapezoid:	area	=	[(a + b) x h] ÷ 2	
Circle:	area	=	radius (r) ² x 3.1416 (π)	= diameter (d) ² x 0.7854
	radius	=	d ÷ 2	
	diameter = circumference	=	r x 2	= π x d
Sphere:	volume	=	r ³ x 4.1888	= d ³ x 0.5236
Triangle:	area	=	(W x H) ÷ 2	
Cylinder:	volume	=	r ² x 3.1416 x L	
<u>Finding Tank Capacity (gallons):</u>				
Cylindrical tanks:	(inches)	=	L x d ² x 0.0034	
	(feet)	=	L x d ² x 5.875	
Rectangle tanks:	(inches)	=	L x W x height x 0.004329	
	(feet)	=	L x W x height x 7.48	
Elliptical tanks:	(inches)	=	L x short diameter (sd) x long diameter (ld) x 0.0034	
	(feet)	=	L x sd x ld x 5.875	




Circle



Square




Trapezoid




Parallelogram




Rectangle



Triangle



Cylindrical Tank



Rectangular Tank

Metric System Conversion Factors

J. Bryan Unruh

Extension Turfgrass Specialist

Area Equivalents

1 acre = 43,560 ft² = 4840 yd² = 0.4047 hectares = 160 rods² = 4047 m² = 0.0016 sq. mile
 1 acre-inch = 102.8 m³ = 27,154 gal = 3630 ft³
 1 hectare (ha) = 10,000 m² = 100 are = 2.471 acres = 107,639 ft²
 1 cubic foot (ft³) = 1728 in³ = 0.037 yd³ = 0.02832 m³ = 28,320 cm³
 1 cubic yard (yd³) = 27 ft³ = 0.765 m³
 1 square foot (ft²) = 144 in² = 929.03 cm² = 0.09290 m²
 1 square yard (yd²) = 9 ft² = 0.836 m²

Liquid Equivalents

1 ft³ of water = 7.5 gal = 62.4 lbs. = 28.3 liters
 1 acre-inch of water = 27,154 gal = 3630 ft³
 1 liter (l) = 2.113 pts. = 1000 ml = 1.057 qts. = 33.8 fl.oz. = 0.26 gal
 1 US gallon = 4 qt. = 8 pt. = 16 cups = 128 fl.oz. = 8.337 lbs of water = 3.785 L = 3785 ml = 231 in³ = 256 tbsp. = 0.1337 ft³
 1 quart = 0.9463 liters = 2 pt. = 32 fl. oz. = 4 cups = 64 tablespoons (tbsp.) = 57.75 in³ = 0.25 gal = 946.4 ml
 1 pint = 16 fl. oz. = 2 cups = 473.2 ml = 32 level tablespoons = 0.125 gal = 0.5 qt
 1 cup = 8 fl. oz. = ½ pt. = 16 tablespoons = 236.6 ml
 1 tablespoon = 14.8 ml = 3 teaspoons (tsp.) = 0.5 fl.oz.
 1 teaspoon = 4.93 ml = 0.1667 fl. oz. = 80 drops
 1 US fluid ounce = 29.57 ml = 2 tablespoons = 6 tsp. = 0.03125 qt
 1 milliliter (ml) = 1 cm³ = 0.34 fl.oz. = 0.002 pts

Temperature Equivalents

degrees Centigrade = (°F-32)x5/9
 degrees Fahrenheit = (°Cx9/5)+32

Pressure Equivalents

1 lb per square inch (PSI) = 6.9 kilopascal (kPa)

Length Equivalents

centimeter (cm) = 0.3937 inch = 0.01 m = 0.03281 ft.
 meter (m) = 3.28 feet = 39.4 inches = 100 cm = 1.094 yds = 1000 mm
 kilometer = 0.621 statute mile = 1000 meters = 100,000 cm = 3281 ft = 39,370 in.
 inch = 2.54 cm = 25.4 mm = 0.0254 m = 0.08333 ft.
 foot = 0.3048 meters = 30.48 cm = 12 inches
 yard = 0.9144 meters = 3 feet = 36 inches = 91.44 cm
 statute mile = 1760 yards = 5280 feet = 1.61 kilometers = 1609 meters

Mixture Ratios

1 mg/g = 1000 ppm
 1 fl.oz./gal = 7490 ppm
 1 fl.oz./100 gal = 75 ppm
 1 pt/100 gal = 1 teaspoons/1 gal
 1 qt/100 gal = 2 tablespoons/1 gal

Flow

1 gpm = 0.134 ft³/minute
 1 ft³/min. (cfm) = 449 gal/hr. (gph) = 7.481 gal/min.

Weight Equivalents

1 ton (US) = 2000 lb = 0.907 metric tons = 907.2 kg
 1 metric ton = 10⁶ g = 1000 kg = 2205 lb
 1 lb = 16 oz = 453.6 grams (g) = 0.4536 kg
 1 oz (weight) = 28.35 g = 0.0625 lb
 1 gram = 1000 mg = 0.0353 oz = 0.001 kg = 0.002205 lb
 milligrams (mg) = 0.001 grams
 1 kilogram (kg) = 1000 grams = 35.3 oz = 2.205 lbs
 microgram (µg) = 10⁻⁶ grams = 0.001 mg
 nanogram (ng) = 10⁻⁹ grams = 0.001 micrograms (µg)
 picogram = 10⁻¹² grams
 1 ppm = 0.0001% = 0.013 fl oz in 100 gal = 1 mg/kg = 1 mg/L = 1 µg/g = 0.379 g in 100 gal water = 8.34 x 10⁻⁶ lb/gal = 1 µl/l
 10 ppm = 0.001% = 10 mg/L 100 ppm = 0.01% = 100 mg/L 1000 ppm = 1mg/g = 0.1% = 1000 mg/L
 1 ppb = 1 µg/kg or 1 µg/L or 1 ng/g
 1 ppt = 1 picogram/g
 1 % = 10,000 ppm = 10g/L = 1g/100ml = 10g/kg = 1.33 oz by weight/gal water = 8.34 lbs/100 gal water

Conversion Factors

To Convert	Multiply by	To Obtain
Acres	43,560	Sq. feet
Acres	0.00405	Sq. kilometer
Acres	4047	Sq. meter
Acres	4840	Sq. yards
Acre-feet	325,851	Sq. feet
Acre-feet	43560	Cu. feet
Acre-feet	1233.5	m ³
Bar	14.5	Lb/sq.in.
Bar	1019.7	g/cm ³
Bar	29.53	inches Hg @ 0°C
Bushels (dry)	0.03524	m ²
Centimeters (cm)	0.03281	Feet
Centimeters	0.3937	Inches
Centimeters	0.1094	Yards
Centimeters	0.01	Meters
Centimeters	10	Millimeters (ml)
cm/sec	1.9685	ft/min
cm/sec	0.0223694	MPH
cm ³	0.0610237	inch ³
Cubic feet (ft ³)	0.0283	Cu. meter
Cubic feet	7.4805	Gallons
Cubic feet	1728	Cubic inches
Cubic feet	0.037	Cubic yards
Cup	8	fl oz
Feet (ft)	30.48	Centimeters
Feet	0.3048	Meters
Feet per minute	0.01136	MPH
Foot candle	10.764	Lux
Gallons (gal)	3.785	Liters
Gal	3785	Millimeters
Gal	128	Ounces (liquid)
Gal/acre	9.354	Liters/hectare
Gal/1000 ft ²	4.0746	L/100 m ²
Gal/minute	2.228 x 10 ⁻³	Cubic feet/second
Grams (g)	0.002205	Pounds
Gram	0.035274	oz
Grams per liter	1000	PPM
Grams per liter	10	Percent
Grams/sq.meter	0.00020481	lb/sq.foot
G/cm ³	0.036127	lb/in ³
G/cm ³	62.428	lb/ft ³
Hectares (ha)	2.471	Acres
Inches	2.540	Centimeters
Inches	0.0254	Meters
Inches	25.40	Millimeters
In ²	6.4516	cm ²
In ³	16.3871	cm ³

To Convert	Multiply by	To Obtain
Kilograms (kg)	2.2046	Pounds
Kg/hectare	0.892	Pounds/acre
Kg/ha	0.02048	lb/1000 ft ²
Kg/L	8.3454	lb/gal
Kilometers (Km)	100,000	Centimeters
Kilometers	3281	Feet
Kilometers	1000	Meters
Kilometers	0.6214	Miles
Kilometers	1094	Yards
Km/h	0.62137	MPH
Km/h	54.6807	ft/min
Kilopascals (kPa)	0.145	Pounds/sq.in. (psi)
Liters (l)	0.2642	Gallons
Liters	33.814	Ounces
Liters	2.113	Pints
Liters	1.057	Quarts
L/100 m ²	0.2454	gal/1000 ft ²
Liters/hectare	0.107	Gallons/acre
Meters (m)	3.281	Feet
Meters	39.37	Inches
Meters	1.094	yards
Meters	100	Centimeters
Meters	0.001	Kilometers
Meters	1000	Millimeters
Meters/sec	2.2369	MPH
M ²	10.764	ft ²
M ³	35.3147	ft ³
M ³	1.30795	yd ³
Miles (statute)	160,900	Centimeters
Miles	5280	Feet
Miles	1.609	Kilometers
Miles	1760	Yards
Miles/hour (mph)	1.467	Feet/second
Miles/hour	88	Feet/minute
Miles/hour	1.61	Kilometers/hour
Miles/hour	0.447	meter/second
Milliliters (ml)	0.0338	Ounces (fluid)
Milliliters	0.0002642	Gallons
Millimeters (mm)	0.03937	Inches
Ounces (fluid)	0.02957	Liters
Ounces (fluid)	29.573	Milliliters
Ounces (weight)	28.35	Grams
Parts per million (ppm)	2.719	lb ai/acre foot of water
PPM	0.001	Grams/l
PPM	8.34	Lb/million gal
PPM	1	mg/kg
PPM	0.013	Ounces/100 gal of water
PPM	0.3295	Gal/acre-foot of water
PPM	8.345	lbs/million gal of water
Percent (%)	10	g/kg
Pint	0.473	liter
pt/A	1.1692	L/ha

To Convert	Multiply by	To Obtain
Pounds	0.4536	Kilograms
Pounds	453.6	Grams
Pounds/acre	1.12	Kg/hectare
Pounds/A	0.02296	lb/1000 ft ²
Pounds/sq.ft.	4883	Grams/sq.meter
Pounds/1000 ft ²	43.5597	lb/A
Pounds/yd ³	0.0005937	G/cm ³
Pounds/gallon	0.12	Kg/liter
PSI (lbs/sq.in.)	6.9	Kilopascals
PSI	0.06895	Bar
PSI	0.068046	atm
Quarts	0.9463	Liters
Qt/A	2.3385	L/ha
Sq. centimeters	0.001076	Sq. feet
Sq. centimeters	0.1550	Sq. inches
Sq. feet	929	Sq. centimeters
Sq. feet	0.0929	Sq. meters
Sq. feet	9.294 x 10 ⁻⁶	Hectares
Sq. inch	6.452	Sq. centimeters
Ton (2000 lbs)	907	kg
Yards	91.44	Centimeters
Yards	0.9144	Meters
Yards	914.4	Millimeters
yd ³	27	ft ³
yd ³	0.7645	m ³
P ₂ O ₅	0.437	P
K ₂ O	0.830	K
CaO	0.715	Ca
MgO	0.602	Mg

Decimal and Millimeter Length Equivalents

Fraction (inch)	Decimals (inch)	Millimeters
1	1.00	25.4
15/16	0.9375	23.812
7/8	0.875	22.225
13/16	0.8125	20.638
¾	0.75	19.05
11/16	0.6875	17.462
5/8	0.625	15.875
9/16	0.5625	14.288
½	0.5	12.70
7/16	0.4375	11.112
3/8	0.3750	9.525
11/32	0.34375	8.731
5/16	0.3125	7.938
9/32	0.28125	7.144
¼	0.25	6.350
15/64	0.234375	5.953
7/32	0.21875	5.556
13/64	0.203125	5.159
3/16	0.1875	4.762
11/64	0.171875	4.366
5/32	0.15625	3.969
9/64	0.140625	3.572
⅛	0.1250	3.175
7/64	0.109375	2.778
3/32	0.09375	2.381
5/64	0.078125	1.984
1/16	0.0625	1.588
3/64	0.046875	1.191
1/32	0.03125	0.794
1/64	0.015625	0.397

Slopes

10% =	6° =	10:1	33% =	18° =	3:1
18% =	10° =	6:1	50% =	26° =	2:1
25% =	14° =	4:1	100% =	45° =	1:1

PESTICIDE APPLICATION RECORD

Company Name _____ Commercial Applicator _____
 Application Date & Time _____ Site Location _____
 Pesticide License Category _____ Number _____
 Pesticide Name(s) _____ Manufacturer _____
 EPA Registration No. _____ Restricted-entry Interval (REI) _____
 Active Material & Formulation _____
 Lot No. _____ % Concentration _____
 Safety Equipment Needed _____

APPLICATION INFORMATION

Type of Area Treated _____ Target Site _____
 Target Pest(s) _____ Total Treated Area _____
 Application Rate (e.g., per acre or per 1000 sq. ft.) _____ Application Timing _____
 Amount of Pesticide Product Mixed _____ Per _____ Gallons of Water _____
 Additives (Surfactant/Wetting Agent/Crop Oil, etc.) _____ Rate _____

WEATHER CONDITIONS

Air Temperature (°F) _____ % Relative Humidity _____ Dew Presence (Y/N) _____
 Initial Wind Velocity (MPH) _____ Wind Direction _____
 First Hour _____ Second Hour _____ Third Hour _____
 Soil Temperature at 4 inches (F) _____ Soil Moisture _____ % Cloud Cover _____

APPLICATION EQUIPMENT

Method of Application _____ Speed (mph) _____ Motor Speed (RPM) _____ Nozzle Type _____ Number _____
 Nozzle Height _____ Spacing Boom Width _____ Gallon Per Acre (GPA) _____ Spray Pressure (PSI) _____
 Nontarget Plant, Animal, or Human Exposure: Yes _ _ No_ _ (If yes, list corrective or emergency action taken)

Other Comments:

Signature _____

_ Date _____

Emergency Pesticide Information

Fred Fishel, Ph.D.
UF/IFAS Pesticide Information Coordinator

Poison Information Center: 1-800-222-1222

The Poison Information Center toll free hotline automatically links a caller in Florida to emergency services on poison prevention and management provided by one of three centers located in Jacksonville, Miami, and Tampa. Each center in the Florida Poison Information Center Network is certified by the American Association of Poison Control Centers as a Regional Poison Control Center and is located on the campus of a major teaching hospital. Emergency and information calls placed to the Network are assessed, triaged, managed and followed by specially trained nurses, pharmacists, physician assistants, physicians and on-call board certified toxicologists.

Web address: <http://www.fpicn.org/>

National Pesticide Information Center (NPIC): 1-800-858-7378

The National Pesticide Information Center (NPIC) is a cooperative effort of Oregon State University and the U.S. Environmental Protection Agency. NPIC is a toll-free telephone service that provides pesticide information to any caller in the United States, Puerto Rico, or the Virgin Islands. NPIC provides objective, science-based information about a wide variety of pesticide-related subjects, including:

- pesticide products
- recognition and management of pesticide poisoning
- toxicology
- environmental chemistry

NPIC staff have toxicology and environmental chemistry education and training to provide knowledgeable answers to pesticide questions. NPIC's toll free call center is staffed 9:30 am to 7:30 pm Eastern time, 7 days a week excluding holidays.

Web address: <http://npic.orst.edu>

CHEMTREC® (Chemical Transportation Emergency Center) 1-800-424-9300

The Chemical Transportation Emergency Center (CHEMTREC®), located in the Washington, DC area, is maintained by the American Chemistry Council. Its purpose is to be a public service hotline for fire fighters, law enforcement, and other emergency responders to obtain information and assistance for emergency incidents involving chemicals and hazardous materials. In addition CHEMTREC helps shippers of hazardous materials comply with the US Department of Transportation Hazardous Materials regulations. Because many companies use CHEMTREC, the emergency number appears frequently on shipping documents, material safety data sheets, rail cars, trucks, and other containers. Companies that list CHEMTREC's emergency number must be registered with CHEMTREC, which includes payment of an annual fee. Thousands of manufacturers and shippers worldwide rely on the CHEMTREC Emergency Call Center to provide information and technical assistance for emergencies involving their products. CHEMTREC maintains a state-of-the-art communications center and a high-end MSDS document storage and retrieval system, containing nearly 2.8 million MSDSs. These sheets are updated and maintained in cooperation with the registrants and are indexed for rapid retrieval.

Web address: <http://www.chemtrec.com/chemtrec/>

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Millie Ferrer, Interim Dean

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition.